

Madras Agricultural Journal

(ORGAN OF THE M. A. S. UNION)

Vol. XXII.]

DECEMBER 1934.

[No. 12.

CONTENTS

	PAGE	PAGE
Editorial	415	
ORIGINAL ARTICLES:		
1. Studies in Paspalum Scorbiculatum, L. The Kodo Millet	419	Rao Sahib V. Muthuswamy Ayyar 446
2. Hill Villages of Tiruppattur Taluk	425	Research Notes 448
3. Hymenopterous Parasites of economic importance in South India	430	Abstracts 449
		Gleanings 452
		Crop and Trade Reports 454
		College News and Notes 455
		Weather Review 456

Editorial.

Livestock in Health and in Disease. "What is the good of concentrating on strong reins to drive a horse or on rubber-tyred wheels to make the carriage comfortable if the horse is so weak that he cannot pull the carriage, or the carriage so frail that it falls to pieces directly anybody sits in it." So said our Viceroy, Lord Irwin, than whom the Indian farmer had no warmer sympathizer. These words were uttered to drive home the importance of animal husbandry to Indian national well-being.

In 1931 Madras had nearly 47 million human beings. In 1930 there were about 16½ million oxen, 13 million sheep, 7½ million goats, 6 million buffaloes, 130 thousand donkeys and 40 thousand horses, moving round about this human throng. Add to this, innumerable mules, pigs, geese, poultry, turkeys and various other domestic animals and it becomes unrealisable in what a crowd of animal beings the human being is enmeshed. A goose lives for 80 years, horse 35, buffalo 25, cattle and pigs 20, goat 15, hen 14, and sheep for 10 years. Many lives of these livestock are thus in a concurrent run with a generation of human lives and as such are in-eliminable factors in the life of man and his well-being.

Dumb as these livestock are, and living as they do lives nearer nature, they have been more the subjects of exploitation in health, than objects of attention in disease. Veterinary science is thus a development later than that of cattle breeding. When man conquered nature, cleared forests, domesticated animals and depended on their support for his well-being, he laid the foundations of the above sciences, that are this day contributing so much to human happiness. They signalize man's efforts to get over the dangers of his existence on earth as a dominant entity. The problems of animal health and welfare are thus intertwined with those of man and his well-being. The position is poignantly so with the agriculturist.

The millions of our cattle, some of them of the finest of breeds in the world, do not all represent a business proposition. The waste of expenditure incurred in India on useless cattle was computed to be something in the neighbourhood of 60 crores of rupees, an amount about equal to that spent on the Army. In any other country the unfit animals would have been slaughtered, but in India where of the triple utility, work, milk and meat, the last is the least, such a course is not easily possible. Mr. Smith has urged that the Indian cultivator cannot even afford to have both buffaloes and cattle, the former for milk and the latter for work. He urged that it will be more economic to have cattle alone and breed them for the dual purpose of work and milk. Against this ideal, the experiments in cross-breeding with foreign blood runs quite counter and has elicited the recent well merited verdict "that the results hitherto achieved in the crossing of indigenous breeds with foreign breeds indicate that this system of improving livestock in India does not produce results of a permanent nature and is not suited to the ordinary condition of the country." The grading up of local breeds is thus imminent and we urge with all the persuasion at our command the revival of the cattle breeding station for Ongoles—a breed that has a world wide reputation.

Concurrently with this, efforts should be made to mitigate the multiplication of our useless animals and the elimination of the drain that they are, in the feeding and upkeep of the good ones. The practice of dedicating scrap animals to the temple, which later on turn out to be the breeding bulls of the village should be modified and with an organization like the Hindu Religious Endowment Board, it should be possible to get into a working arrangement by which, such dedication is confined to good pedigree calves supplied by cattle farms. All animals of dubious parentage should be castrated as they are thus replaced. The action of the Punjab Government in empowering District Boards to frame regulations for the castration of scrub animals could usefully be followed by us.

The efforts of the State at cattle improvement are not enough. In England gentlemen farmers have taken the lead in cattle breeding.

They have thus shown a considerable spirit of service to the farming community and to the State. The State by itself cannot do all that is indicated as necessary in this direction. In Madras except the isolated instance of the Pattagar of Palayakottai, there are no noteworthy endeavours in this direction. It is time the fact was brought home to our landed magnates that their wealth and opportunities are merely a trust which has both privileges and responsibilities and that one of the most potent ways in which they could function towards their country and discharge their debt to their bretheren in agriculture is to provide them with good breeding bulls that will enrich their stock of work and milk animals.

Livestock are a remarkable vehicle for the conversion of earth's vegetative growth into food that has become indispensable to man. Meat, milk and eggs are some of these. Our civilization has become complex and the production and distribution of these involve organization, centralization and hygenic supervision. In the side of medicine, humanity is becoming more and more dependant on glands and other parts of animals for the treatment of diseases. The manufacture of anti-toxins and serums needs organized production and control, so that animal products can be standardised and be made safe and dependable.

Diseases due to malnutrition are as common among animals as among men. Rotational grazing is recommended. But above all, researches into grasses have to be initiated and a systematic attempt to handle herbage should be made. With a proper handling of the nutritional aspects of animal health, much of the troubles in connection with animal disease will be successfully tackled.

Diseases that man has to fear through animal infection are anthrax, rabies, tuberculosis, and innumerable other parasitic infections. The degree to which these are controlled among animals will provide the measure by which he could successfully ward them off from his ambit. It is therefore becoming imperative that the maintenance of animal health is a *sine qua non* to that of keeping up human health.

Veterinary education is centred in urban areas. It loses that benefit which comes to it with the intimate touch with livestock which rural areas give. It has therefore been urged that to mitigate this want of intimate contact, model dairies should be attached to Veterinary Colleges. The initiating of riding classes has also been recommended. These measures connote antidotes to the tendency of a mere academic contact with animals and in a measure rectify a vital defect. The need for an Institute of Veterinary Research for our province is keenly felt and it is our hope that parallel to agriculture and in proximity to it, there will soon be established, a centre of Veterinary Research.

A new orientation in our outlook is indicated as a necessity. The usual view of looking at veterinary service as merely an instrument of preventive and curative medicine should cease. The problem of disease should be viewed from the bigger back-ground of Animal Health. The maintenance of animal efficiency in health should be the major issue. What the future Veterinary Officer should be is best surmised in the words of Sir Daniel Hall " We want to see a class of men who have charge of a district, who are thinking about the horses, the cattle, the sheep and the pigs and how to keep disease away from them. Naturally, they will have to know about the endemic diseases, but breeding, environment, nutrition, and other factors in hygiene will be equally important." Such a concept may be very long before it is realized in India; but the creating of a post of Rural Commissioner that we have advocated will hasten this consummation.

Rao Sahib V. Muthuswami Ayyar. The end of 1934 sees the termination of the active official service of M. R. Ry. Rao Sahib V. Muthuswami Ayya Garu, Lecturer in Agriculture, at the Agricultural College. His wide and varied experience of agricultural matters gave him a great hold on all that came in contact with him. His readiness to serve them found in him a guide, philosopher and friend. A gentleman of high principles, of a very sympathetic disposition, cast in a mould of service, Mr. Muthuswami Ayyar will ever remain a unique personality to his friends and students. His very active habits have not abated with age and we hope and pray that it will be given to him to have the satisfaction of utilising the rest of his life to the service of the poor and the depressed. His charitableness is patent. We pray that it will please God to keep him for many years to come, in good health and vigour, for the service of the Motherland.

STUDIES IN *PASPALUM SCROBICULATUM*, L. THE KODO MILLET

By G. N. RANGASWAMI AYYANGAR, B.A., I.A. S.,
Millets Specialist, Agricultural Research Institute,
AND V. PANDURANGA RAO, M.A.,
Assistant, Millets Breeding Station, Coimbatore.

The Kodo millet, *Varagu* (Tamil) and *Arika* (Telugu), occupies an area of well over a million acres in the Madras Presidency. The grain of this millet is easily preserved and proves a good famine reserve. It is a poor man's food. The crop is very drought-resistant. Only well-matured grains should be husked and used as food. Husks and immature grains are poisonous. This millet is considered safer for use as the grains get old. Both as food and fodder the crop is poor. While the crop is ripening it is considered dangerous to graze animals on it. This millet is grown mostly rain-fed, though in small areas under irrigation also. Trichinopoly and South Arcot districts have over 170,000 acres each; Kurnool, Nellore and Ramnad have each over 90,000 acres. The Nilgiris, Malabar and South Kanara are the only districts in which this crop is not grown. About 21,000 acres are raised under irrigation. Of these Coimbatore, Ramnad, Madura and Kurnool have about 13 thousand acres. This millet yields on an average 800 lb. of grain per acre with a range of 400 to 1200 lb. according to the tract. The yield of straw is about 1000 to 2000 lb. per acre. The straw is used as a manure in alkaline lands. This millet has been under study at the Millets Breeding Station and the following summarise the knowledge so far gained.

Seedlings. The seeds of this millet have a thick husk which is about 40 per cent. of the seed by weight. It is, therefore, slow in germination. On the fifth day after sowing are seen the first signs of sprouting. A single long leaf is visible and no stem is in evidence. All leaf-sheaths and under-surfaces of leaves are densely hairy. The mesocotyl is smooth with no rootlets and dries up soon. It is capable of elongation adjusting to varying depths of sowing. It is unpigmented though the adult plant develops purple pigment. When the seedlings are a fortnight old, the first secondary root makes its appearance. Secondary roots are also unpigmented.

Adult Plants. These are usually erect, and occasionally spreading or prostrate in habit and are 1 to 2½ feet in height. They differ in their leafiness. In a good leafy plant the leaves are broad and numerous and give it a dense, bushy appearance. Less leafy plants have narrow leaves giving them an all-stem and little-leaf look. The nodes are swollen or not according to varieties. The first node is hairy and the other nodes are glabrous. In a few varieties villous upper

nodes are met with. The internodes are solid. The length of internodes increases gradually from bottom to top in any tiller. The internodes, in most cases, are fully ensheathed. The number of tillers in varieties varies from 5 to 18. The early tillers set well. The later ones are more sterile. Nodes touching the ground strike root. They ascend from a prostrate and rooting many-noded base; simple or sparingly branched, usually five-noded. The leaf blades are linear to lanceolate-linear; equally wide or slightly constricted at base; glabrous; margins scabrid; smooth on either surface, broad or narrow; erect, arched or bent. The arched condition is associated with the broad leaf which droops and gives the plant its characteristic appearance. Narrow-leaved plants have very few bent leaves. The leaf is usually of a dark green colour. A few varieties have light green leaves. The midrib of the leaf is white or dull green in colour. The ligule is membranous, colourless, broad or narrow. The junction of the leaf blade and sheath is usually covered with a felt of long hairs. Rarely such hairs are short. The leaf tip remains green in most varieties but in some it dries up to brown even before the heads ripen. The size of leaf varies with the varieties and a considerable range in their length has been noted resulting in differential habits.

Purple Pigmentation. This manifests in seedlings as a purple wash. With the growth of the seedling and the elongation of the leaf-blade, leaves begin to bend and from the bend to the tip there is a wash of purple. At a further stage of growth when the basal nodes show out, their nodal bands take on the pigmentation. With increased growth, such bits of internode as the intense ensheathing leaves bare, begin to have a wash of purple more of self colour and less of lines. Leaf junctions, axils and leaf margins are purple coloured. In some the tips of glumes develop purple. The stigmas show grades of purple ranging from dark to very faint purple. In the anthers purple rings and dots are developed on a yellow background. Anthers without any easily noticeable purple pigment occur occasionally. The stigmas and anthers dry to a brown colour of grades which parallel the depth of purple in them. After the flowering is over, the panicle branches get liberated from the sheaths and the peduncles also get coloured purple. The optimum manifestation of the purple pigment is from the flowering to the milky stage of the grain, after which there is a falling off. Advancing age, however, loosens out the internodes from the clasp of the sheaths and these together with prominent nodal bands retain the purple pigment as long as they continue to be sappy. Looked at *en masse* this field crop has a characteristic violet look. Though all the varieties are purple pigmented, the depth of colour varies. In some the colour is so much reduced that only a very careful examination reveals the pigment in odd parts.

Panicle—Emergence. The panicle arises usually from the fourth node. The first sign of the growing panicle is a slight bulge in the

leaf-sheaths. The swelling increases gradually and the panicle is seen through the sheath slit. It takes about a week for it to emerge. When fully emerged it is enclosed by three overlapping leaf sheaths, which inroll one inside another at the tops. The flag and the leaf below it, likewise inroll at the top and, in most cases, do not separate even when the panicle is in full flower. In the inrolled region mentioned above the ends of the inflorescence remain stuck up. The upper end thus fixed, the floral branches, as they elongate, bend outwards at the centre. The enclosing leaf-sheaths are forced apart by the rapid growth and consequent arching of the panicle until thereby the tips are also forced out. This is the general rule but in a variety imported from Sierra Leone, the peduncles clear out of the leaf-sheaths, elongate and hold the panicles aloft and away from the leaf-sheaths.

Panicle—Arrangement. Excepting the lower three nodes, the other two bear panicles. From each of the two upper nodes three separate peduncles arise. Of these the central one aborts and the two lateral grow. The abortive one bears a sessile panicle. This panicle may contain one or two branches bearing rudimentary spikelets. Of the two free growing peduncles one grows quickly and shows out earlier than the other. The earlier is always bigger and has more spikelets and less of sterility. In the later peduncle the spikelets at the tips of the branches are usually sterile. Each of these peduncles gives rise to three branches one of which aborts, the other two showing a likewise differential growth and development. These branches repeat this trimerous process until small spikelet-bearing branches of differential size arise and produce flowers. In this ultimate trimerousness some of the abortives get converted into long stalked single flowers. The weaker flower-bearing branches are mostly unbranched, whereas the stronger ones branch out and bear a larger number of flowers. The largest number of spikelets in a branch may be as many as 100.

Panicle—Branches. Each branch has a broad, flat rachis with a series of depressions corresponding to the situation of the spikelets. On the side on which the spikelets are situated, a central ridge runs along the entire length. On either side of the ridge the spikelets are arranged alternately in two series on short pedicels. In some varieties instead of the usual two-seriate condition, a branching of the pedicel gives rise to a non-seriateness, the spikelets being irregularly arranged. At the base of the branch it is two-seriate; in the middle the non-seriate condition prevails; and towards the tip the two-seriateness continues. The non-seriate condition is found in some cases along the entire length of the branch. Other variants to the simple branching of the pedicel and the production of two flowers are the following:—(1) the pedicel instead of forking into two, branches into three, each bearing a spikelet. (2) It may bear more than four flowers at different

levels. (3) Small branches arise at different levels on the ridge of the branch and these bear a fairly large number of flowers. In addition to the crowding induced by the branching of the pedicel, the double seededness of such non-seriate heads is the most important factor in the crowding and small size of the grain in the earheads that are not two-seriate. The disturbance of two-seriateness brings about a dense packing of spikelets on the flat rachis and contributes to the reduction in the size of the spikelets. Consequently the spikelets in these are $\frac{1}{3}$ to $\frac{1}{2}$ the size of those in the two-seriate panicles. It may be observed that the non-seriate varieties are early, lighter pigmented with grains of a lighter brown husk.

Spikelet—Single Seeded. The description of this spikelet has been elaborated from Hooker's Flora of British India in which a description of the variety with double seeded spikelets is not found. Spikelets orbicular, mostly decidedly plano-convex, falling entire from the short rudimentary pedicels and abaxial on the dilated rachis of spike-like racemes.

Glume I. o (suppressed.)

Glume II. More or less equal to the spikelet; convex; membranous; light green; deciduous; glabrous; 5-6 nerved.

Glume III. Similar to Glume II, but less convex and more flat; light green; thin; glabrous; deciduous; 2-5 nerved; along the inner margins are seen shallow transverse pits whence the specific name "*scrobiculatum*."

Glume IV. Horny; pale green; later develops a light or dark brown colour; glabrous; 5 transparent nerves; margin firm; obtuse; emucronate; persistent.

Palea. Tightly embraced by the narrowly involute margins of Glume IV; similar in substance to Glume IV; 2 transparent nerves; the palea with flaps widened into a broad auricle below the middle; persistent.

Stamens. Three; filaments short, 1 m. m. long, anthers-3, 2-3 m. m. long; 2-loculed; locules open by longitudinal lateral sutures

Ovary. Oval; translucent; stigmas-2, styles distinct and laterally exerted near the tip of the floret; styles feathery from one-third the length from the apex.

Lodicules. Two; fleshy; serrated tips; broadly cuneate.

Grain. Tightly enclosed by the slightly hardened glume and palea; rotundate-elliptic; very convex in front, flat on the back; pale; scutellum up to half the length of the grain.

Spikelet—Double Seeded. In the double-seeded spikelet, between Glume II and palea of Glume IV an extra flower is interpolated. It is enclosed in an extra glume and palea. This extra flower develops seed, each spikelet thus having two seeds. Abortive conditions of this double seededness freely occur interspersed with this double fertility and arise as follows. An extra flower is developed between Glume II and palea of Glume IV. It is a perfect flower and has only an extra palea. It does not set seed.

Opening of the Flower. The opening of the first flower is generally on the second day after the emergence of the panicle. This

flowering begins between 2-30 and 3-0 a. m. on each day and continues till sunrise. Youngman and Roy (1923) note that these flowers open between 7-30 and 8-0 a. m. at Nagpur. Flowers do not commence to open from any definite region. They usually start from the middle of the floral branch and gradually spread to either ends. Instances in which flowers begin to open at either end are met with occasionally.

Anthesis of a Flower (anthers extruded). Detailed observations on the anthesis in a normal average flower are recorded below showing the trend of sequence.

2-30 a. m.	Glumes begin to open.
2-40 "	Anthers visible through opening.
3-15 "	Anthers emerge.
3-30 "	Anthers completely out.
3-35 "	Anthers dehisce.
3-45 "	Glumes close completely.

The stigmas may or may not come out of the glumes. When the glumes begin to gape the anthers crowd at the orifice and are undehisced. They are mostly non-emergent. Their filaments are 1 mm. in length. In stray cases anthers emerge, their filaments being 6 mm. long. These filaments remain turgid for a long time, often till 8-9 a. m. The anthers may emerge simultaneously or one by one or two at a time followed by the third. This stray emergence accounts for the paucity of evidences of flowering in this unobtrusive millet. When the anthers remain inside the flower, their dehiscence takes place long after the opening of the glumes. Dehiscence starts as a slit at one end and gradually spreads or it begins in the middle and proceeds to the ends. The stigmatic feathers dry in the evening. The anthers remain fresh and do not wither till next morning. The lodicules are fleshy and do not shrink immediately after the anthesis of flowers but remain fleshy for 6--8 hours after the opening of the glumes and then dry up—a probable device preventing the closing glumes from jamming the anthers.

Progress of Flowering. The following table connotes the daily anthesis energy of the 15 per cent of flowers opening during the flowering period, emerging and non-emerging anthers included.

Day of Flowering.	2-3 a. m.	3-4 a. m.	4-5 a. m.	5-6 a. m.	Total.
First Day	14	1	1	-	16
Second Day	2	10	-	1	13
Third Day	7	11	-	-	18
Total.	23	22	1	1	47

N. B. 15 per cent of the flowers in the head opened. The remaining 85 per cent were cleistogamous.

It will be seen that all the flowers in the head do not open. The percentage of open flowers in the varieties varies from nil to 50 per cent., the most frequent being 10 to 15 per cent. This millet is

therefore highly cleistogamous which explains the complete absence of natural crossing in it. Any artificial manipulation of the spikelet irretrievably damages it and many attempts at emasculation and artificial pollination proved futile.

Grain. The grain matures in 30–35 days after flowering. It is tightly enclosed by the hardened fourth glume and its palea. The husk is coloured shades of brown. In the two seriate varieties, the grain is bigger and nearly twice the size of those in the non-seriate ones. In the variety imported from Sierra Leone (referred to above), though the head is two-seriate, the grain is small. The degree of emergence of the heads has no effect on the setting of the seed, both having about the same degree of sterility. In this millet seed-setting is dependent on the season. In the year 1928, the drought affected the seed-setting in some varieties to such an extent that not even a few grains per plant could be obtained. Under favourable conditions good yields can be expected.

False Polyembryony. In the year 1931, in the course of a number of seed germinations for albinism in this millet, two instances of two seedlings arising from a single seed were noticed. The seedling had a single root and two plumules each with its own coleoptile. At the surface of the seed, the two were separate. Paraffin sections of the seed showed that the plumules had independent vascular bundles. Lower down, the cortical portions of the two shoots were found to unite, the bundles running separate. In a few sections lower down, the vascular bundles were closer and approached each other until finally they became enclosed in a single endodermis. Sections still lower down showed the root strand run into this single bundle. From this examination of the course of the vascular bundles and also of the cortical region, it will be seen that what appears to be independent at the top is, in reality, the result of the branching of a single seedling. At a very early stage in the development, the mesocotyl has branched into two, resulting in the double seedling—a case of false polyembryony. Cases of pseudo-polyembryony have been recorded by other workers. Only those pertaining to the Gramineae are noticed. In maize, Kiesselbach (1926) noted seedlings with (1) two plumules each with its own coleoptile and two primary roots enclosed in a single coleorhiza, (2) a single plumule with two primary roots in a single coleorhiza. In the case reported here, there are two plumules each with its own coleoptile but with a single radicle.

Paspalum Sanguinale, Lamk. Several species of *Paspalum*, especially *P. dilatatum* are grown in America, Australia and South Africa as pasture grass. A wild ally of *Paspalum* known as *Chicco* (*P. sanguinale*, Lamk) is grown in the Vizagapatam District. This was grown at the Millets Breeding Station for a number of years and the following notes are appended.

The seed which is very small takes 7 days to germinate, two days later than for *Varagu*. Unlike *P. scrobiculatum*, the seedlings are green and have no purple pigment anywhere in the plant. Before flowering the plants are spreading and almost prostrate. The panicle-bearing tillers become erect at flowering. Unlike *Varagu* the internodes are hollow and much exposed. At ripening stage they have a golden yellow colour. The nodes are glabrous and not swollen. The flag is the broadest leaf in the plant. The upper surface of leaves is rough and the lower is smooth. The leaves are arched and not bent. The plants have five to six heads with long well emerged straight peduncles. Occasionally their fullness leads to goose-necking. The panicle has a general resemblance to that of a well grown *Chloris barbata*. It consists of a number of fingers (or branches) arranged in irregular whorls along a short axis. An average earhead may have about 40 fingers. The bulk of these arise from the two bottom whorls, the rest of them being distributed to those above, mostly in twos and threes. Each finger may have about 100 flowers. The spikelets are arranged alternately in clusters of one to three. The spikelets are very small, the length in each being four to five times the width. The glumes are prominently ribbed and dry to a straw colour. The structure of the spikelet is like any other *Paspalum*. The flowers of this wild ally open from 1–30 a. m. and the anthesis continues up to 7 a. m. The greatest anthesis energy is within the first hour after opening. It takes four to five days for a panicle to complete its flowering.

This wild *Paspalum* scores over *P. Scrobiculatum* in a number of points, viz., more herbage, free earheads, greater drought resistance, larger number of seeds per head and absence of sterility. A cross with this wild ally is indicated as a potential source of improving the Kodo millet, if the difficulties in the manipulation of this close and delicate cleistogamous flower could be overcome.

References.

1. Kiesselbach, T. A. (1926). False Polyembryony in Maize. *American Jour. Bot.* 13. 33–34.
2. Youngmann, W. and Roy, S. C. (1923). Pollination Methods among the lesser millets. *Agr. Jour. India.* 18: 580–583.

HILL VILLAGES OF TIRUPATTUR TALUK.

By U. B. MOHAMED ABBAS, B. Sc. Ag.

Agricultural Demonstrator, Tirupattur, N. A.

Introduction. A portion of the Eastern Ghats is situated in Tirupattur Taluk, N. Arcot district in two separate blocks viz. the Yelegiris and the Javadhi hills. The entire length of the mountain range is 45 miles in the taluk. The Yelegiri has a range of 13 miles and is situated a mile and a half from Jalarpet Railway Station, while the Javadhis with a range of 32 miles run continuously from north to

south forming a boundary on the eastern side of the taluk. The height of the hills above the sea level ranges from 2,700 to 3,900 feet. The total extent of the mountain range can be classed as (a) Reserved Forests 33,641 acres, (b) Patta lands 10,924 acres, (c) Grazing grounds 22,713 acres. *Kambugudi* the highest plateau in the Javadhi range is 3,900 feet above sea level and is 24 miles from Tirupattur town. There is a travellers' bungalow in the middle of the plateau with a good motorable road leading to it.

There are 43 villages in all, 12 in the Yelegiris and 31 in the Javadhi with a total population of 14,245 according to the latest census. Each village has 15 to 50 houses and most of them can only be reached by foot paths while a few have bridle paths. There are a few travellers' bungalows scattered here and there on the top of the Javadhi and the Yelegiri hills. The scenery from the top of the hills is very picturesque. Some parts of the hills are covered with dense vegetation and wild animals like bison, sambar, cheeta, bears and deer are found.

Climate. Owing to the favourable height of the mountain range the climatic conditions on the top of the hills are very good. The temperature ranges between 60 and 90°. The rainfall is also not very heavy, being only about 35 inches per annum and major portion of this is received during the north-east monsoon season (October to December).

People and their habits. The inhabitants of these hill villages belong to aboriginal tribes known as *Malai-yalis*. Recently a few non-malai-yali villages have also sprung up in favoured localities, by migration of the cultivating classes of the neighbouring plains. The *Malai-yalis* call themselves as Gownders and have no sub-castes. They worship both Vishnu and Siva. Each village has a Kali or a Dhurga temple. Vishnu and Siva temples are few and are held in common for all the neighbouring villages. The requirements in life of these primitive people are very few. There is no shop in any of these villages. The few things they require are salt, kerosene, matches, betel leaves, nuts, mud pots and clothes. Clothes they purchase once a year at the time of Pongal, one of the two festivals they celebrate. The rest they obtain from the neighbouring weekly shandies of the plains. The barber, black-smith, carpenter and other artisans come from the plains at regular intervals in the busy season, complete their work in the village and return to the plains. Every village has a family of Adi-Dravadas of the plains whose services are utilised during marriages, festivals and funerals and in return they are paid in kind annually.

A group of villages have a headman in common and he with the help of a Panchayat, a committee to which every house of the village contributes a member, settles the disputes and the fines and penalties collected go to the village common good fund which is utilised for the

celebration of Pongal or Kali festival. The people have two festivals viz. Pongal celebrated for ten days in the month of January and the other Kali or Dhurga festival celebrated in the summer months. Marriages are also regarded as public functions since all the villagers are related to one another and so every one takes part in the function. The marriage expenses are met by the bridegroom or his party and may cost about Rs. 100 besides 15 to 20 bags of paddy or ragi. Some of the villages are very malarial but the original inhabitants are highly immune to it.

There are three village Panchayat Elementary Schools for all the 43 hill villages. The one on the Yelegiri is due to the efforts of a few Anglo-Indian families that have settled there and the other two are for the children of Reddiyur and Naickenoor villages on the Javadhi inhabited by people migrated from the neighbouring plains. It will thus be seen that there is not even a single school for the *Malai-yali* villages. The postman visits some of the villages once a week and the staff of the Health Department twice a year.

Occupation. The people depend mainly on agriculture and partly on cooly work they do to the Forest Department. An average ryot has 5 to 10 acres of dry land and there is very little of wet land. There are very few wells fit for irrigation purposes, so the people have little or no chance of raising a crop except with the help of rain-fall. Major portion of the annual rain-fall is received in the north-east monsoon season i.e. October to December which is the main agricultural season. The soil on the slopes and the plateau is either red loam or sandy loam with a good mixture of kankar or sandstone to a depth of 6 inches or even up to one foot with soft stone and rock underlying it. The soil has therefore very little retentive power but responds well to heavy manuring.

Ragi, cumbu, samai, varagu, tenai, cholam, castor, groundnut, mustard and wheat are raised as rainfed crops. Besides these crops, some ryots have a few plantain and jack trees. Wherever there is facility for irrigation from jungle streams, or small tanks, paddy and sugarcane are raised. In the southern portions of the Javadhi hills near Nellivassal, coffee is grown under the shade of jack trees and it fetches Rs. 50 to 150 from about 10 cents (about 100 plants) of land. With the advent of the south-west monsoon, the fields are prepared and if the season happens to be good, cumbu mixed with cholam is sown and red gram dibbled in lines. Other millets except ragi are also sown in June—July. Groundnut is also sown at the same time. In August, after a good rain, ragi is sown and castor dibbled. A few ryots sow short duration paddy also. In September—October, wheat and mustard are sown. The rain-fed crops are intercultivated only once with a country plough.

From September onwards harvest of crops begins. Cumbu is the first crop to come off the field while castor comes last. Coffee plants flower in May—June and the berries are harvested in December—January. Ryots who have wells fit for irrigation, cultivate paddy thrice in the year. Besides paddy, cholam, ragi, sugar cane, are also grown under irrigation. Every ryot has a few heads of cattle and they form a source of income. During the period the crops are on the field, the ryots graze their livestock in the forests for which they pay 8 annas per cow or bullock per annum.

The probable income of a ryot who has 10 acres of dry land with a pair of work animals, 10 calves and 6 jack trees with 5 members in the family is shown in the *appendix*. In an average year a ryot may get Rs. 10 as net profit. If he happens to be thrifty, he may save some money, acquire land from the Government, clear the forest and bring it under cultivation. To purchase and clear the forest and bring it under cultivation, it costs him about Rs. 40 per 10 acres. The savings are used for the purchase of young calves and not for the purchase of jewels, nor does he waste it on drink. Money lenders were rare and the hill tribes never used to borrow as a rule but nowadays they come down to the neighbouring plains and borrow.

Except for the two forest roads which touch a few villages, the others have to be reached only by means of bridle paths which are very rugged. They are so bad in some places that even the pack animals find it difficult to negotiate them. Hence the ryot never brings down his bulky produce like ragi and cumbu to the neighbouring shandies for sale, but has to get his groundnut, mustard, jack fruit and coffee. The hill tribes are badly in need of a fair weather road, medical aid and elementary education.

Conclusion. The following are a few agricultural improvements that can be effected with success:—

- a) distribution of improved strains of crops grown there e. g. E. C. 593, *Surangi* cumbu, Adt. 3 paddy etc.
- b) introduction of light iron ploughs of cooper 25 type.
- c) demonstration of better methods of cultivating crops e. g. wheat and coffee.
- d) introduction of money crops e. g. potato and pine apple.
- e) inducing ryots to grow fruit trees e. g. mangoes, oranges, pomegranates, jack, and cashew nuts. These trees come up very well.
- f) stationing breeding bulls with the village Panchayats for the improvement of hill cattle.

The writer's thanks are due to Mr. M. Kanti Raj, Assistant Director of Agriculture, Vellore for the encouragement and suggestions given in preparing this article.

APPENDIX

Income in kind and cash.

1. Ragi	4 acres	500	Madras measures
2. Cumbu	2 acres	200	"
3. Groundnuts	3 acres	1,800	"
4. Castor dibbled with ragi	...	200	"
5. Dry paddy	½ acre	200	"
6. Red gram dibbled with cumbu		200	"
7. In the remaining portion he has his chillies (dry) plantains, mustard and other things which he uses for himself.			
8. By the sale of 5 heads of cattle @ Rs. 12 each		Rs. 60	0-0
9. By the sale of groundnuts (1800 m. m.) @ Rs. 3/- 200 m. m.		...	Rs. 27-0-0
10. By the sale of Jack fruits		...	Rs. 6-0-0
11. Do. Do. Castor		...	Rs. 4-0-0
12. By engaging himself or some member of the family as cooly to the forest department.		Rs.	3-0-0
			<u>Total Rs. 100-0-0</u>

Expenditure in kind and cash.

1. Ragi	500 m. m.	8 months		Rs.	30-0-0
2. Cumbu	200	3	"	Rs.	6-0-0
3. Paddy	200	1½	"	Rs.	10-0-0
4. In the months of April and May when jack fruits are available, raw fruits are boiled and consumed with ragi gruel.				Rs.	5-0-0
5. Cost of clothes	Rs.	30-0-0
6. Grazing charges	Rs.	6-0-0
7. Cost of 5 young cows for rearing	Rs.	10-0-0
8. Kist at 8 annas per acre	Rs.	5-0-0
9. Cost of kerosene, matches, salt, betel leaves, nuts, mud vessels each at 8 as. a week.				Rs.	26-0-0
10. Festivals Pongal and Kali Puja	Rs.	10-0-0
11. Purchase of plough parts sickles etc.	Rs.	2-0-0
12. Marriages to be attended	Rs.	1-0-0
				<u>Total Rs. 90-0-0</u>	

Net income is Rs. 10-0-0

HYMENOPTEROUS PARASITES OF ECONOMIC IMPORTANCE IN S. INDIA

BY T. V. RAMAKRISHNA AYYAR, B.A., Ph. D., F. Z. S.

&

V. MARGABANDU, M.A.

(*Madras Agricultural Department*)

The study of economic entomology has in these days attained such dimensions that various methods of control are being advocated for checking insects injurious to crops. The mechanical methods of control such as the use of insecticides, spraying, fumigation and other artificial devices such as the use of light traps, etc., have been in some cases found impracticable and uneconomical. In view of the poisonous nature of the insecticides themselves, the prohibitive cost of the treatments in some cases, the frequency and thoroughness with which these treatments have to be given, the promptness with which these have to be done, and the several meteorological factors that influence the success or failure of the results obtained and lastly the temporary and palliative nature of the results obtained, mechanical methods of control frequently become very much limited in their scope.

During recent years a new method of crop-pest control has sprung up known as the biological control of insect pests. It is a well known fact that almost every insect has at one stage or another, its parasite or predator. A parasite is one which becomes attached to the host, at some stage of its existence, and leads an ecto or endozoic existence gradually killing the host; whereas a predator is one which leads a free existence and preys on its victim killing the same immediately in most cases. Among insects, the phenomenon of parasitism is very highly developed in some of the groups of Hymenoptera, especially the Ichneumonoid families, Braconidae, Ichneumonidae, Chalcidae and Bethyilidae and the Tachinids among the Diptera in the insect world.

The object of this paper is to record the different kinds of hymenopterous parasites so far found in the South Indian region possessing some economic importance and to set forth instances, where some of the insect hosts are controlled, to a greater or lesser extent by the natural enemies. A study of this was undertaken in order to get an idea as to the indigenous parasites that exist in the region with their respective hosts, the parasite or parasites that attack a particular host, the degree to which there is parasitism and the regions where they are obtained. This would help us in getting an idea as to the possibilities for breeding parasites and liberating them in numbers to control crop pests.

A list of hymenopterous parasites with their respective hosts has been given at the end noting their distribution. Work in these parasitic groups was begun in S. India by the senior author and lists were prepared by him, one in 1919 (2), a second one in 1921, (3) and the most recent one in the Bulletin of Entomological Research, (10). There are stray publications such as the Ann. & Mag. Nat. History., Proc. U. S. Nat. Mus., Jl. Bom Nat. Hist. Socy., Bull. Ent. Res., etc. wherein S. Indian parasites have been described by such writers as Rohwer, Gahan, Girault, Crawford, Cameron, Silvestri, Grandi, Waterston, Wilkinson, Ferriere and a host of others.

Instances where crop pests are being controlled by natural enemies noted in South India are:

Nephantis serinopa, the black headed caterpillar is a very serious pest of coconut palms both in the East Coast as well as in the West Coast. And this is a classic example wherein natural enemies are used to a degree to control the pest. In the East Coast the pest is not serious probably on account of the presence of the parasites which have a favourable environment for successful multiplication; whereas in the West Coast probably on account of the secondary parasites the primary parasites have not asserted themselves. Hence breeding laboratories were opened at three different stations, viz., Calicut, Mangalore and Ponnani where the parasites were bred in large numbers and liberated. This pest has got parasites attacking the larval and the pupal stages. There is a Bethyloid, a Microbracon, an Apanteles and an Elasmid attacking the different stages of the larva; and a Chalcid, Eulophid and an Ichneumonid attacking the pupa in addition to a Tachinid fly. In spite of so many parasites attacking a single host at different stages of the pest, it has been found difficult to control the pest on account of the secondary parasites on every one of these. The coconut caterpillar complex has been well shown diagrammatically (11a).

The life history and habits of the larval parasite *Elasmus nephantidis*, R., has been worked out at greater detail (15).

In the case of the paddy stem borer *Schoenobius incertellus* there are the egg parasites as well as the larval parasites. The chief egg parasites are—(1) *Trichogramma minutum*, R., (2) *Tetrastichus schoenobii*, Ferr. and (3) *Phanurus beneficiens*, Zehnt. Of these *Tetrastichus schoenobii* would appear to be most effective in that a single grub is able to kill 2 or 3 eggs of the host. These parasites are found during the months of February and March during the *dalva* or the second crop and this is the reason why the infestation is very low during this crop and the infestation very high during the *salva* or the first crop in the Circars. In addition to these there are the Braconid larval parasites *Tropobracon indicus*, R., *Microbracon* sp., and *Apanteles schoenobii* and the Ichneumonids *Goryphus maculipennis* and *Ischnojoppa luteolator*, F. But instances of these asserting themselves are not known.

Argyria sticticraspis. Hmps. The sugarcane borer is another in whose case in other parts of the world, biological methods of control are being carried on by utilising the egg parasite *Trichogramma minutum*, R. In India, so far the damage done to cane has not warranted a control of the pest in this direction. The egg parasite *Trichogramma minutum*, R. and *Phanurus* sp. are found in the Circars and among the larval parasite, *Stenobracon* sp.

Sesamia inferens. Wlk. the ragi borer is a clear instance where the parasites have proved of great utility under natural conditions. The infestation is high only in the *pairu* or cold weather crop and here the larval parasites check the pest. A braconid wasp keeps the pest under control. They are noticed from February to March and the percentage of parasitisation goes on increasing from February onwards till the percentage rises to 80% during the first week of March. Here is a case where the parasites have been found really useful in checking the pest at the right time unlike the egg parasites of the paddy stem borer.

Utetheisa pulchella. Linn. on Sannhemp: Egg parasites probably *Phanurus* sp. noticed on the eggs are very prominent and the parasitisation is very high during the second week of March. Eggs were collected at different periods to note the degree of parasitisation which went on rising from February to March. It was 57.24% during the last week of February; 63.82% during the first week of March and 87.88% during the 2nd week of March. In addition to these there are Tachinids parasitic on the larvae which are also high during these months; and a chalcid hyper-parasite was noticed on these Tachinid maggots.

Leaf miners on citrus: *Phyllocnistes citrella* Stn. Chalcid parasites were noticed on the pupae. Though the miners were prominent right through the year, the parasites were noticed only from February up to April. The percentage of parasitisation was worked out as 33.15. The parasites were not to be seen during the remaining part of the year.

The eggs of the Fulgorid *Pyrilla perpusilla* Wlk. which is found on cane shows parasitisation by Chalcids Dryinids. *Pyrilla* in large numbers on the cane leaves cause sooty mould on account of the secretion of honey dew and the whole crop looks black and may be detrimental to the crop too; and the breeding of egg parasites and liberating them may be tried, this depending upon the degree to which the pest attacks the crop.

The cholam borer—*Chilo simplex* Butl. The list at the end would show that this has got Ichneumonids, a number of Braconids and a Chalcid parasite on it.

There are a number of scale insects and mealy bugs infesting many trees and plants and every one of these is attacked by a parasite or

another. Such serious scales as the *Pulvinaria maxima*. Green. on nim, *Ceroplastodes cajani* on *Zizyphus jujuba*, *Aspidiotus orientalis*. Newst. on tamarind, *Lecanium* species on cotton, sandal, etc., are a few instances where the parasites check them. The hymenopterous wasps especially of the super family Chalcidoidea are parasitic on the many species of Coccids. These lead an endozoic existence at least in the early stages and lead a free life in the adult stage. Parasitisation by these minute insects are not visible until after the adult parasites emerge when the exit-holes could be seen. In certain cases where the scales are transparent there will be abnormal colour. Study in these cases is rather difficult. A study of the parasites of Coccids would form a study by itself.

There are the several important crop pests of the Noctuid family often subject to parasites such as *Spodoptera mauritia*, Boisd. and *Cirphis albistigma* that attack paddy, *Prodenia litura* F.B. the tobacco caterpillar, *Achaea janata*, Linn. a serious pest on castor, *Eublemma olivaceae* Wlk. the brinjal leaf roller; and among the Pyralids several of the paddy pests, sugarcane pests, cotton and vegetable pests, have also parasites. The list at the end will give an idea as to the range of orders and families in which we find insect parasites. But for the existence of these parasites, all the crops would have been overrun by these pests and there will be no crops at all. Nature in her bounty and generosity is maintaining a balance, as it were, even without our consciousness, thus restricting the scope of these pests. This will give one an idea as to the utility of these little insects to man.

General considerations. The general life history of these parasitic wasps may be stated thus. The adult is a free living insect. It goes in search of the right host and lays an egg or eggs. The eggs may be laid singly or in an eggmass on or in the body of a caterpillar, maggot or grub. The parasites are provided with an apparatus at the tail end called the ovipositor which is used for inserting the eggs. They are even so adapted that these can be thrust from outside into the body of larvae inside the plant stems. The hatched out larvae may lead an ectozoic or endozoic existence and feed upon the tissues and vital organs of the host bringing about the gradual death of the host. The grubs after feeding on the host may come out, and spin cocoons on the surface of the host very often as in the case of Braconid wasps.

These minute wasps which we find in thousands are widely different in structure and habit and a study of these would bring to light many interesting points which would help us in the efficient utility of these parasites toward insect pest control. These may be ectozoic or endozoic and most of these are parasitic only in their larval stages. The list at the end will show that a particular species of parasite may have different hosts; a number of species may attack a single host simultaneously; some may be polyphagous; and some restricted only to particular hosts. In the case of the black scale

Saissetia nigra in California there are 51 species recorded on it. The original parasites are in some cases attacked by secondary parasites and this phenomenon is called hyper-parasitism. Some attack insects belonging to different orders or families or even groups and different stages of the hosts. We have species of *Apanteles* attacking larvae of different orders. These parasites attack the different stages of the host. There are egg parasites, larval parasites, and pupal parasites. Thus it could be seen that the host may be attacked at any stage or in all stages. There is the Trichogrammid—*Trichogramma minutum*, R., attaching the eggmasses of *Schoenobius*, *Argyria* and *Diatraea*. There is *Phanurus beneficients*, Zehnt, attacking the eggmasses of *Schoenobius* and *Scirpophaga*. Similarly there are instances of larval parasites. Many of the Braconids and Ichneumonids are larval parasites. Among the pupal parasites there is the Chalcid *Pluristropis epilachneae*, Roh attacking the pupa of *Epilachna*, *Tetrastichus ayyari*, *Chilo simplex*, *Acanthojappa* sp., *Melunitis ismene* and so on. Among the hyper parasites we have *Perilampus microgastris*, Ferr., on *Microgastris indicus*, Wik., *Apanteles machaeralis*, Wik., and a Braconid parasite on *Nepantis serinopa*, M., *Mesochorus plusciphilus*, V., on *Apanteles plusiae*, Vier, and *Marietta* sp. on some of the parasites on Coccids. Specialisation has gone to such a degree that many of the Eurytomids are phytophagous and the Chalcid family *Agaonidae* live on seeds of figs. Members of the family Eurytomidae attack galls. There is the seed chalcid *Bruchophagus mellipes*, Gahan, boring into *Sesbania* seeds.

In certain cases where we notice more than one species attacking a host and where the parasites lead an endozoic life it requires a more careful and thorough study as we should be able to distinguish between the primaries and secondaries. There are instances of parasitism wherein they are parasitic on the predators of the host themselves in which case they are not beneficial. There is *Aphrastobracon flavipennis*, Ashm. which is parasitic upon *Eublemma* which is a predator on Coccids; similarly there is *Elasmus indicus* found with the Coccid *Anomalococcus indicus* which is probably predaceous on *Eublemma*. This clearly shows that a clear understanding of the interrelations of the host and the parasite are necessary.

Thus we could see how varied are the nature and kind of parasitisation, the habits of these parasites and the relationship between them and the hosts. A study of the natural enemies available in South India brings to light the following interesting points. Parasites are not found throughout the season but are found only during particular seasons, e. g., in the case of *Schoenobius* we have the egg parasites from January to March during the summer months at Samalkota; similarly we find the larval parasites from January to March in the case of the ragi borer and the egg and larval parasites of Sannhemp caterpillar and the Chalcid parasites of the citrus leaf miner could be

had only during the months of January to March. Here we find some interesting features. One is that parasites are found only during particular seasons; in these cases cited they are found only during the months of February—March; in some cases they are useful in controlling the pest as in the case of Ragi borer; in some cases though the crop exists during two seasons and the pest also is found the parasite could be found in abundance only during the second crop as in the case of *Schoenobius* and similarly in the case of *Utethesi* on Sannhemp and the citrus leaf miner *Phyllocnistes*. In the case of *Nephantis* in the West coast it becomes serious in certain seasons but the pest becomes controlled as soon as the parasites assert themselves. Thus it could be seen that there are certain meteorological and other factors that now and then put the parasites under check.

Before initiating "Biological methods of control" it is incumbent upon us to know before-hand the indigenous parasites that exist in the area with the respective hosts; and their host relations; the number and kind of parasites that attack a particular host and the number of hosts which a particular species attacks; the secondary parasites that may be found on the original parasites; the different stages of hosts that are attacked by the parasites; the seasonal abundance of these parasites and their numerical relation with that of the host; and the factor or factors that are responsible for the success or failure for the multiplication of these. All these points require elucidation. Then one should be able to eliminate the adverse factors and attempt at successful breeding of the parasites and liberating them, the kind of cages to be utilised depending upon the nature of parasites—whether egg or larval.

The list of parasites given below will show how vast the field is and how there is plenty of scope for utilising the parasites in the control of crop pests. In view of the complexity of this method of control it requires a careful study before any one starts on the work of biological control; and in this connection one may be referred to such eminent men like Howard, Thompson, Smith, Burgess, Jones and a host of others who are doing a good deal of work on insect parasites and biological methods of control. A good deal of literature by these men has accumulated and would prove very interesting study in this new line of work.

The list does not by any means profess to be complete but it is just given to show the richness of parasites found on the several major crop pests thereby showing one that there is a vast and wide field of study of great economic importance as well as of deep academic interest. It may also be added that the list gives only those South Indian forms of economic importance and does not deal with the numerous parasitic forms which affect other insects in various ways.

List of Hymenopterous Parasites of Economic Importance noted in South India.

Host family.	Host insect.	Parasitic insect.	Parasite family.	Distribution.
LEPIDOPTERA				
Hesperiadae	<i>Parnara mathias</i> (Rice skipper) do. do. do. (on pupa)	<i>Xanthopimpla immaculata</i> , Mor. <i>Clinocentrus</i> sp. <i>Eupsteromatus parnaeae</i> , Gah. <i>Ischnojoppa luteator</i> , F.	Ichneumonidae Braconidae Pteromalidae Ichneumonidae	Palur (South Arcot), Madras & Coimbatore. Nellore, Karvetnagar. Coimbatore.
Lasiocampidae	<i>Taragama sica</i> (on Acacia) <i>Taragama dorsalis</i> <i>Lasiocampid larva</i> <i>Trabala vishnu</i>	<i>Hentcospilus reticulatus</i> , Cam. <i>Apanteles taragamae</i> , Vier. <i>Chalcis argentifrons</i> , Ash. <i>Microdus fumipennis</i> , Bing.	do. Braconidae Chalcididae Braconidae	Coimbatore. Bangalore. Coimbatore.
Limacodidae	<i>Parasa lepida</i> (on castor) do. do. & <i>Thoses</i> sp. <i>Conthelya rotunda</i> (on coconut palm) <i>Natada nararia</i> (on <i>Pithecolobium</i>)	<i>Clinocentrus</i> sp. <i>Stomatocerus ayyari</i> , Gah. <i>Eurytoma parasae</i> , Gah. <i>Alciodes</i> sp. <i>Protapanteles</i> sp.	do. Chalcididae Eurytomidae Braconidae do.	Coimbatore. Coimbatore. Coimbatore. Malabar. Coimbatore.
Lymantridae	<i>Olene mendosa</i> <i>Euproctus fratarna</i> (on castor) do. (rose leaves) <i>Euproctus scintillans</i> and <i>E. fraternus</i> (on larva) <i>Euproctus scintillans</i> (on gogu)	<i>Goryphus</i> sp. <i>Hentcospilus (reticulatus?)</i> <i>Disophrys</i> sp. <i>Hentcospilus merdarius</i> , Gr. <i>Protapanteles</i> sp.	Ichneumonidae do. Braconidae Ichneumonidae Braconidae	Coimbatore. Coimbatore. Madras. Saidapet and Coimbatore. Coimbatore.

<p><i>Orgyia postica</i> (on larva) <i>Psalis securis</i> (on paddy)</p>	<p><i>Apanteles (Protapanteles) colemani</i>, Vier. <i>Protapanteles</i> sp.</p>	<p>do. do.</p>	<p>Mysore. Coimbatore.</p>
<p><i>Polyptychus dentatus</i> (on Cardia) (egg parasite) <i>Daphnis nerii</i> (on larva) <i>Macroglossum</i> sp. (an undescribed species) (on larva)</p>	<p><i>Anastatus coimbatorensis</i>, Gir. <i>Troporhogas maculipennis</i>, Cami. <i>Microplitis</i> sp.</p>	<p>Chalcididae Braconidae do.</p>	<p>Coimbatore. Coimbatore. Coimbatore.</p>
<p><i>Achaea janata</i>, Linn. (on larva) do. do. do. do. (on castor) do.</p>	<p><i>Tetrastichus ophiusae</i>, Craw. <i>Microtorfidea lissonata</i>, Vier. <i>Zamesochorus orientalis</i>, Vier. <i>Microplitis</i> sp. <i>Paniscus ocellaris</i>, Th. <i>Microplitis ensyrus</i>, Lyb.</p>	<p>Eulophidae. Ichneumonidae. do. Braconidae. Ichneumonidae. Braconidae.</p>	<p>Mysore. Mysore. do. S. India. Coimbatore. All over S. India. ("a specific and effective parasite on the castor semi-looper") Coimbatore.</p>
<p>do. do. <i>Prodenia litura</i> (on larva) do. do. <i>Plusia peponis</i> do.</p>	<p><i>Euplectrus leucostomus</i>, Robl. <i>Hemicosphilus</i> sp. <i>Apanteles prodeniae</i>, Vier. <i>Diocetes argenteopilosa</i>, Cami. <i>Chalonus</i> or <i>Chaloualia</i> sp. <i>Apanteles plusiae</i>, Vier. <i>Mesochorus plusiophilus</i>, Vier. (a hyper parasite on <i>Plusia</i>) <i>Paracopidosomopsis joravae</i>, Gir. <i>Ceraphron aethanasi</i>, Gir.</p>	<p>Eulophidae. Ichneumonidae. Braconidae. Ichneumonidae. Braconidae. do. Ichneumonidae Encyrtidae. Proctotrypidae.</p>	<p>Coimbatore. do. Mysore. Coimbatore. do. Mysore. Bangalore. Coimbatore. do.</p>
<p><i>Plusia signata</i> <i>Plusia agramma</i> (on snake gourd) <i>Carea subtilis</i> (on <i>Eugenia</i>) <i>Adisura abhinasoni</i> (on lab-lab)</p>	<p><i>Tumidicoxoidae jambilana</i>, Gir. <i>Microbracon</i> sp.</p>	<p>Chalcididae. Braconidae.</p>	<p>do. do.</p>

Host family.	Host insect.	Parasitic insect.	Parasite family.	Distribution.
LEPIDOPTERA (Contd.)				
Noctuidae (Contd.)	<i>Cirphis</i> sp. <i>Cirphis abtistigma</i> (on paddy) <i>Spodoptera mauritia</i> do. do. (on barley) (on pupa). <i>Earias fabia</i> <i>Perigææ cepensis</i> do. (infesting saf- flower plants) do. <i>Eublemna olivaceæ</i> (Brinjal leaf roller) do. <i>Eublemna scitula</i> A Noctuid larva (on maize stalk)	<i>Meteorus</i> sp. <i>Xanthopimpla</i> sp. <i>Chelonus</i> sp. <i>Charops</i> sp. <i>Ichneumon</i> sp. <i>Microbracon lefroyi</i> , Dudge. & Gough. <i>Euplectrus eplerias</i> , Roh. <i>Heterogamus percurrans</i> , Lyle. <i>Protapanteles</i> sp. <i>Microdus</i> sp. <i>Hymenobosmena</i> sp. <i>Aphrastobracon flavipennis</i> , Ash. <i>Cremastus noxiosus</i> , Mori. <i>Xanthopimpla pedator</i> , F. <i>Iphiaulax</i> sp. <i>Merionotus</i> sp. <i>Apanteles flavipes</i> , Cam. <i>Apanteles (Stenoplectura)</i> sp. <i>Iphiaulax spilocephalus</i> , Cam. <i>Glyptomorpha (Bracon) deesææ</i> , Cam. <i>Apanteles</i> sp. (probably new). <i>Microbracon ciliocida Ramakrishna</i> .	Braconidae Ichneumonidae Braconidae Ichneumonidae do. Braconidae Eulophidae Braconidae do. do. Ichneumonidae Braconidae Ichneumonidae do. do. do. do. do. do. do. do. do.	South Arcot. Palur (S. Arcot). Malabar. South India. Ootacamund. Ramnad. Coimbatore. do. do. do. do. S. India. Coimbatore. do. Coimbatore, Kurnool. Coimbatore. Poona. Godavari & Coimbatore S. India. do. Mandya. S. India.
Pyrilidae	<i>Chilo simplex</i> (on Sorghum) do. do. do. do. do. do. do.		Braconidae do.	do. Coimbatore, Kurnool. Coimbatore. Poona. Godavari & Coimbatore S. India. do. Mandya. S. India.

do.	<i>Xanthopimpla nrsai</i> .	Ichneumonidae	All over India.
do. (on pupa)	<i>Tetrastichus ayyari</i> , Roh.	Eulophidae.	Coimbatore.
Schoenobius (eggmass)	<i>Tetrastichus schoenobii</i> , F.	do.	Hebbal, Mysore.
do.	<i>Phanurus beneficiens</i> , Zehnt.	Scelionidae.	do.
do.	<i>Trichogramma minutum</i> , R.	Trichogrammidae.	do.
do. (larva)	<i>Tropobracon indicus</i> , R.	Braconidae.	do.
do.	<i>Microbracon</i> sp.	do.	do.
do.	<i>Goryphus maculipennis</i> , Cam.	Ichneumonidae.	Godaveri Dt.
do.	<i>Apanteles schoenobii</i> , W.	Braconidae.	Hebbal, Mysore.
do.	<i>Ischnojoppa tuteator</i> , F.	Ichneumonidae.	Very common in Coimbatore & all S. India.
Scirpophaga (eggmass)	<i>Phanurus beneficiens</i> , Zehnt.	Scelionidae	Bangalore.
Diatraea (eggmass)	<i>Trichogramma minutum</i> , R.	Trichogrammidae.	Hebbal, Mysore.
<i>Sylepta derogata</i> (cotton leaf roller)	<i>Neopimploides sylepta</i> , Vier.	Ichneumonidae	Mysore.
do.	<i>Microtoridea lissonata</i> , Vier.	do.	do.
do.	<i>Elasmus indicus</i> , Roh.	Elasmidae.	S. India.
<i>Euzophera perticella</i>	<i>Pristomerus euzopherae</i> , Vier.	Ichneumonidae.	Bangalore.
<i>Leucinodes orbonalis</i> (brinjal fruit borer)	<i>Pristomerus testaceus</i> , Mort.	do.	Attur, Chinglepet.
<i>Euzophera perticella</i>	<i>Phanerotonia</i> sp.	Braconidae.	Coimbatore.
<i>Pyrausta nacheaeratis</i>	<i>Apanteles machaeratis</i> , Wlk.	do.	Nilambur, Madras.
<i>Cnaphalocrocis medanales</i> (Paddy leaf-roller)	<i>Cardiochilus</i> sp.	do.	Ganjam.
<i>Etiella zinckenella</i> (Infesting sunnhemp)	<i>Phanerotonia</i> sp.	do.	Coimbatore.
<i>Crocizelonia binotalis</i> (on Radish)	<i>Microbracon</i> sp.	do.	do.
do. (pest of cruciferous plants)	<i>Microbracon lefrovi</i> , Dud & Gough.	do.	S. India.
do.	<i>Microbracon mellatus</i> Ramakrishna.	do.	do.

Host family.	Host insect.	Parasitic insect.	Parasite family.	Distribution.
LEPIDOPTERA (Contd.)				
Pyrilidae (Contd.)	<i>Antigastra catalaunatis</i> (Gingelly leaf-caterpillar) <i>Dichocrocis</i> (on castor seed) <i>Pyralid larvæ</i> (boring into the fruit & pods of <i>Pongamia glabra</i>) <i>Scirpophaga</i> (egg-mass)	<i>Hymenobosmana</i> sp. <i>Diocetes trochanterata</i> , Mordl. <i>Microbracon pictus</i> , Ramakrishna.	Ichneumonidae. do. Braconidae.	Coimbatore. do. do.
Pterophoridae	<i>Diatraea</i> (egg-mass) <i>Euselastis atomasa</i> (on Red-gram)	<i>Phanurus beneficiens</i> , Zehnt. <i>Trichogramma minutum</i> , R. <i>Protapanteles</i> sp.	Scelionidae. Trichogrammidae. Braconidae.	(Bangalore) Hebbal, Mysore. Hebbal, Mysore.
Gelechiidae	<i>Stomopteryxnerteria</i> <i>Platyedra</i> (cotton Bollworm) do. <i>Pthorimaea blapsigona</i> (on brinjal bud)	<i>Chelonella</i> sp. <i>Microbracon lefroyi</i> , Dudgeon & Gough. <i>Chelonus</i> or <i>chelonella</i> sp. <i>Microbracon incarnatus</i> , Ramakrishna.	do. do. do. do.	South Arcot. India. South India. do.
Xyloryctidae	<i>Nephandia serinopa</i> do. do. (on pupa) do.	<i>Elaemus nephandiæ</i> , Roh. <i>Stomatocerus sulcatusculetum</i> , Gri. <i>Trichospilus pupivora</i> , Ferr. <i>Xanthopimpla punctata</i> , F.	Elasmidae. Chalcididae Eulophidae. Ichneumonidae	do. do. Cochin. Calicut.
Gracillariidae	<i>Cyphosticha coenuta</i> <i>Gracillaria soyella</i> (on Red gram)	<i>Euryscotinax Coimbatorensis</i> Roh. <i>Anipsiella indica</i> , Gir.	Eulophidae. Chalcididae.	Coimbatore. India.
Hypsidae	<i>Nyctemera lactinia</i> (on larva)	<i>Euplectrus nyctemeræ</i> , Craw.	Eulophidae.	Bangalore.

Arctiade	<i>Hypsa ficus</i> (on <i>Ficus indicus</i>) <i>Larva of?</i> <i>Crentonotus albistriga</i>	<i>Disophrys</i> sp. <i>Meteorus arctiicida</i> , Vier. <i>Apanteles</i> (Protapanteles) <i>creatonoti</i> , Vier.	Braconidae.	Coimbatore. Mysore. do.
Papilionidae	<i>Diacrisia obliqua confusa</i> Hairy caterpillars. <i>Faptilio demoleus</i> & P. <i>polytes</i> . <i>Stauropus alternus</i> <i>Phycodes radiata</i>	<i>Apanteles obliquae</i> , Wlk. <i>Henicospilus horsfieldi</i> , Cam. <i>Apanteles</i> (Protapanteles) <i>papiliotis</i> Vier. <i>Apanteles</i> (Protapanteles) <i>stauropi</i> , Vier. <i>Apanteles phycodis</i> , Vier.	do. do. do. do. do.	Nilambur, Madras. S. India. Mysore. do. do.
Eucosmidae	<i>Argyroproct illepidata</i> do. <i>Cricula trifenestrata</i> <i>Azygophleps scalaris</i> On different plants <i>Psychid larva</i> <i>Melanitis</i> (on paddy) on pupa <i>Ocinara</i> <i>Vioachola</i> (on soapnut fruit)	<i>Eugathis cryptophlebiae</i> , Vier. <i>Trissomalus fulvicornis</i> , Roh. <i>Xanthopimpla pedator</i> , F. <i>Pristomerus</i> sp. <i>Phanarotoma</i> sp. <i>Goryphus nursei</i> , Cam. <i>Acanthojoppa</i> sp. <i>Goryphus?</i> <i>fuscinerus</i> , Cam. <i>Charops obtusus</i> , Mort.	do. do. do. do. do. do. do.	do. do. do. do. do. do. do.
Nymphalidae				
Bombycidae				
Lycaenidae				
COLEOPTERA				
Curculionidae	<i>Alicidas affaber</i> (attacking <i>Hibiscus cannabinus</i> stem borning weevil) <i>Alicidas bibo</i> (on the grub) <i>Actenemis</i> sp. (on the grubs) <i>Calandra oryzae</i> (Rice weevil) <i>Aspidomorpha</i> (on grubs)	<i>Aphrastobracon alcidiphagus</i> , Rank. <i>Campyloneurus ceylonicus</i> , Cam. <i>Ipobracon dentiscapa</i> , Ramakrishna. <i>Pezomaius oryzae</i> , Cam. <i>Tetrastichus colemani</i> , Crav.	Braconidae do. do. Pteromalidae Eulophidae.	Coimbatore. do. do. Salem. . All over S. India. Bangalore.
Cassididae				

Host family.	Host insect.	Parasitic insect.	Parasite family.	Distribution.
COLEOPTERA (Contd.)				
Buprestidae	<i>Buprestid</i> borer <i>Sphenopteri</i> spp. (on larvae boring into groundnut, cotton & pulses)	<i>Vipis gracilis</i> , <i>Ramakrisna</i> . <i>Glyptomorpha (Iphiatulax) smeenus</i> Cam.	Braconidae. do.	Coimbatore. Coimbatore, Bellary.
Ptinidae	<i>Sitodrepa</i> (infesting stored Coriander)	<i>Meraporus vandinsi</i> , Tuck. = <i>Aplastomorpha catandrae</i> , How.	Super. Chalcidoidea.	Madras.
Bruchidae	<i>Bruchus cinensis</i> (on larva) do.	<i>Bruchocida orientalis</i> , Craw. <i>Bruchobius colemani</i> , Craw.	Eupelmidae. Pteromalidae	Bangalore. Bangalore & Coimbatore.
Bostrychidae	<i>Dinoderus</i> (shot-hole borer in bamboo frame work)	<i>Spathius</i> sp.	Braconidae	Coimbatore.
Coccinellidae	<i>Epilachna</i> (grubs) do. (on pupa)	<i>Pleurotropus foveolatus</i> , Craw. <i>Pleurotropus epilachnae</i> , Roh.	Eulophidae. do.	Bangalore. Coimbatore.
Chrysomelidae	Coccinellid grubs <i>Aspidomorpha militaries</i> (on larva)	<i>Homalotylus flaminus</i> (Dalm). <i>Tetrastichus colemani</i> , Craw.	do.	Bangalore.
Cerambycidae	<i>Xylotrechus</i> .	<i>Metapina</i> sp. (probably new.)	Eupelmidae.	do.
DIPTERA				
Anthomyiidae	<i>Cholism</i> maggot	<i>Tetrastichus nyenitawus</i> , Roh.	Eulophidae.	Coimbatore.
Trypetidae	Fruit fly <i>Bactrocera</i> sp. (on Alangium fruits) <i>Carpoomyia vesuviana</i> (attacking Zizyphus fruits) <i>Dacus</i> (<i>Chaetodacus</i>) <i>cucurbitae</i>	<i>Diachasma minorpha comperii</i> , Vier. <i>Bathyaulax trypaeniptaya Ramakrisna</i> . <i>Bathyaulax carpoxyia</i> , Ramkr. <i>Opius fletcheri</i> Sato.	Broconidae. do. do.	India. Anantapur Dt. Coimbatore.
	<i>Dacus longistylus</i> (attacking seed capsules of Calotropis)	<i>Austroopius</i> sp.	do.	Throughout India.
			do.	S. India.

<i>Dacus (Chaetodacus) incisus</i> (infesting fruits of the tree of Careya arborea)	<i>Opus incisus</i> , Sivo.	do.	Coorg.
<i>Dacus (Chaetodacus) incensus</i> .	<i>Opus (Biosteres) persulcatus</i> , Silv.	do.	do.
On a fly boring into Zizyphus fruits	<i>Opus (Biosteres) compansans</i> , Sivo.	do.	do.
On fruit flies	<i>Opus (Biosteres) carponyiae</i> , Silv.	do.	Coimbatore.
On fruit fly maggots (collected and taken to Australia by Compere in 1907).	<i>Asobara orientalis</i> , Vier.	do.	S. India.
<i>Asphondylia sesami</i> (Gingelly gall-fly)	<i>Syniomosphyrum indicum</i> , Sivo.	Super. Chalcidoidea	
Galls on <i>Macarena armeria</i> -probably on gall-flies	<i>Eurytoma dentipectus</i> , Gah.	Eurytomidae.	Coimbatore.
From cumbu grains probably on gall-flies	<i>Tetrastichus coimbatorensis</i> , Roh.	Eulophidae.	do.
From grass galls-probably on gall-flies.	<i>Tetrastichus isaaci</i> , Roh.	do.	do.
<i>Pachydiplosis oryzac-Paddy</i> gall flies,	<i>Tetrastichus okawus</i> , Roh.	do.	do.
<i>Syrphus</i> (on water melon)	<i>Neanastatus trochantericus</i> , Gah.	Encyrtidae.	do.
	<i>Xestonodidea foersteri</i> , Gah.	Platygasteridae	do.
	<i>Folygnotus</i> sp.	do.	Tanjore.
	<i>Bassus orientalis</i> , Cam.	Ichneumonidae.	Coimbatore.
HEMIPTERA			
Aphidae	<i>Aphidius colemani</i> , Vier.	Braconidae.	Bangalore.
	<i>Aphidencyrtus</i> sp.	Encyrtidae	do.
	<i>Leurocerus</i> n. sp.	do.	do.
	<i>Paranagrus optabilis</i> , Perh.	do.	Coimbatore.
Fulgoridae			

<i>Lecanium virida</i> do.	Coccophagus sp. do.	Aphelinidae do.	S. India.
<i>Lecanium discrepans</i> do.	<i>Aphycus</i> sp. Coccophagus sp.	Encyrtidae Aphelinidae	do. do. do.
<i>Lecanium olace</i> <i>Saussurea nigra</i> do.	<i>Aphycus</i> sp. <i>Scutellista cyanea</i> , Mot. <i>Microterys kotinskyi</i> , Fullaway. <i>Encyrtus barbatus</i> , Timb. <i>Scutellista cyanea</i> , Mot.	Encyrtidae Miscogasteridae	Coimbatore. do. do.
<i>Ceroplastes confiferus</i> <i>Inglisia chelonoides</i> <i>Anomalococcus indicus</i> <i>Phenacoccus insolitus</i> <i>Phenacoccus iceryoides</i> <i>Tacharditis lacca</i> <i>Icerya Aegyptiaca</i> <i>Mytilaspis piperis</i> Coccids on <i>Acacia</i>	<i>Scutellista cyanea</i> , chalcid. <i>Elasmus indicus</i> , R. A Eulophid chalcid 3 or 4 small chalcids. <i>Gampylotenicurus indicus</i> , R. 2 small chalcids. A Eulophid chalcid. <i>Elasmus indicus</i> , R. <i>Hadronotus futuiventris</i> . <i>Telenomus colemani</i> , Craw. <i>Anastatus colemani</i> . <i>Telenomus laticulcus</i> .	Encyrtidae Elasmiidae	S. India. do. do. do.
<i>Clavigrella gibbosa</i> (on eggs) <i>Dolycoris indicus</i> (on eggs) <i>Degonetus serratus</i> (on eggs) <i>Coptosoma cribraria</i>		do.	Coimbatore.
Locustid eggs on Cordia leaf <i>Heteroglyphus banian</i> (on eggs) (on paddy) <i>Oxya velox</i> on 'paddy (on eggs) do. do.	<i>Pediobopsis locustivora</i> , Koh. <i>Scotis</i> sp.	Proctotrypidae Platygasteridae Encyrtidae Platygasteridae	Bangalore. Mysore. Bangalore. Coimbatore.
<i>mantis</i> sp. (egg capsules)	<i>Anastatus</i> sp. <i>Tumidiscapus oshlagus</i> , Gir. <i>Podagrion</i> sp.	Eulophidae. Sclionidae. do.	do. do. do.
Mantidae		Super. Chalcidoidea.	
ORTHOPTERA			
Locustidae			

References.

1. Ayyar, Ramakrishna T. V. (1909) Lefroy's "Indian Insect Life"—Hymenoptera section.
2. " (1919) *Proc. 3rd Ent. Meeting, Pusa* Vol. III. pp. 931—936.
3. " (1921) *Proc. 4th Ent. Meeting, Pusa* pp. 29—33, 363—366.
4. " (1921) *J. Bom. N. H. Soc.* Vol. XXVIII. pp. 298—300.
5. " (1923) *Proc. 5th Ent. Meeting, Pusa* pp. 352—362, 263—269.
6. " (1925) *Spolia Zeylonica*, Vol. XIII, Part 2, pp. 235—254.
7. " (1925) *J. Bom. N. H. Soc.* Vol. XXX, pp. 487—491.
8. " (1925) *Departmental Digest for Ju.*
9. " (1926) *Bull. Ent. Res.* Vol. XVII, Part pp. 91—98.
10. " (1927) *Bull. Ent. Res.* Vol. XVIII, Part.
11. " (1928) *Mem. Ind. Dept. Agri. Ent. Ser.* Vol. X, No. 3.
- 11 (a). " (1932) *Madras Agricultural Journal* Vol. XX, No. 4, p. 145.
12. Cherian, M. C. (1928) *Year Book, Madras Agri. Dep.* pp. 12—22.
13. Crawford, J. C. (1913) *Proc. U. S. Nat. Mus.* Vol. 45, pp. 241—260.
14. Gahan, A. B. (1919) *Proc. U. S. Nat. Mus.* Vol. 56, pp. 513—524.
15. Rao, Ramachandra and Cherian, M. C. (1926) *Year Book, Madras Agri. Dept.*
16. Rao, Ramachandra and Cherian, M. C. (1927) *Year Book, Madras Agri. Dept.* pp 11—22.
17. Rohwer, S. A. (1921) *Ann. Mag. Nat. Hist. Ser.* 9, Vol. VII, p. 123.
18. Weld, J. C. (1922) *Proc. U. S. Nat. Mus.* Vol. 61 (includes 3 Indian forms).

RAO SAHEB V. MUTHUSWAMI AYYAR

Mr. V. Muthuswami Ayyar was born on the 16th of December 1879 in the village of Kummalakudi in Tanjore district. His later, intimate, though official, connection with practical agriculture. was more than foreshadowed by his parentage, his father and grandfather being tillers of the soil in their ancestral property of a few acres in their native village. Mr. Muthuswami Ayyar, through his maternal uncle, who lived at Madura, had a number of chances of visiting that historic town and having his early years impregnated with that religious outlook which the associations of the ancient temple served to implant on all impressionable natures in early childhood. Muthuswami's education however was completely confined to St. Peter's College, Tanjore, from where he passed out in 1898. He immediately joined the College of Agriculture in Saidapet, from where he passed out with honours in 1902 winning the Robertson Medal. He was selected by the Department for further training and after a year's stay at Koilpatti, was deputed to the Bombay Presidency where he served on the Poona, the Surat and the Nadiad Farms. On his return, he

served as Farm Manager at Palur, Samalkota, Taliparamba and Anakapalle for various periods up to the year 1918, when he was transferred to Coimbatore as the Teaching Assistant in Agriculture. His long official career of over 32 years is thus defined into two periods—the earlier one from 1902 to 1918, having been spent on some of the biggest farms of this department, and the latter a period of another sixteen years spent at the Agricultural College, Coimbatore.

His choice as teaching assistant was therefore a very happy one, for, he came equipped to his task with the experience gained as a Farm Manager of several stations. Circumstances were less favourable to him than to others of his equipment and training, and it was not till almost the fag end of his career that he was gazetted as Lecturer in Agriculture and that too in an officiating vacancy. Government, however on the eve of his retirement recognised his meritorious service by the award of a Rao Saheb in the birthday honours in June last. He laid down his office on the 16th of December 1934.

Rao Saheb Muthuswami Ayyar in addition to his official duties at the College was intimately and personally connected with a number of activities on the Estate. The Officers' Club, the Officers' Mess, the Co-operative Stores, the Co-operative Society, the Radhakalyanam celebrations, and the Madras Agricultural Students' Union all owe a great deal of their present position to his selfless labours for their inception, or growth or maintenance. The Union especially, and the Journal bear and will bear the stamp of his personality for many more years to come. For a number of years 1920 to 1926 he was an important office-bearer, either as Secretary or as Editor or as Sub Editor or as Vice President and during those five years the responsibility for steering the Union and the Journal was chiefly in his hands, and this he did with great credit drawing unto him by his own example, a band of devoted workers who have helped to build up our traditions.

Rao Saheb Muthuswami Ayyar was a great social worker. He was the friend of the poor, the downtrodden and the despondent. His great heart flowed out in sympathy towards the sufferings of others, and many are those, who when in trouble, domestic or official, have been consoled by his comforting words and guided by his sage advice. He was held in great veneration by all estate residents. By his simplicity—both of dress and of habits—his outspokenness, his accessibility and his readiness to place himself at the service of any one who required him, he endeared himself not only to his students and his colleagues but to his superiors as well.

Rao Sahib Muthuswami Ayyar was an ideal teacher, who combined in himself erudite knowledge and the power to draw out the best from his students. With his quiet unassuming ways and almost paternal

solicitude for his pupils, he created a feeling of reverence in their minds and a desire to be as well informed in practical agriculture as he himself was. He was in fact something like the gurus of old, who not only by precept but also by example, set the standard of high thinking and plain living before their sishyas.

After thirty-two years of active service one cannot grudge him the rest that he so richly deserves; but it is the expressed opinion on many a quarter, that with him, will pass out a mass of information and an encyclopaedic amount of details about agriculture in this Presidency. It is the sincere wish and request of all his students that in the days of his retirement, he will, for the sake of the generations yet to come, translate all his ideas into concrete expositions that will serve as books of reference to the future student of agriculture. Personalities such as his shed light, lustre and love, with peace and good will prevailing, all round them and it is our prayer to the Almighty that Rao Saheb V. Muthuswami Ayyar will be spared very many more years, not only to enjoy his happy retirement in sound and active health, but also to contribute to making his little world of ours with its factions and strifes, a more cheerful and a more worthy place to live in. (M. R. B.)

Research Notes.

The effect of X-rays on Sorghum Pollen.

X-rays are a potential source for inducing mutations. Most of the work on X-radiation is confined to the irradiation of seeds—dry or germinating—under different conditions. In some cases the entire plant is irradiated, while in others the embryo in all stages of development is subjected to X-rays. Very little work with irradiated pollen has been done. The only records available, so far as we are aware, are those of Stadler and McKay and Goodspeed.

Pollen of sorghum was exposed to X-rays for 5, 10 and 15 minutes with a set up of 70 k. v., 1 m. a. target distance of 16 cm. without aluminium filter, and subsequently germinated in artificial media. The germinating medium was—41 per cent sucrose with 1.5 per cent shred agar. It will be seen from the following table that the different dosages have not resulted in statistically significant differences.

Duration of exposure to X-rays.	No. of pollen grains on the medium.		Percentage of germination.
	Sown.	Germinated.	
5 minutes.	1,232	371	30.1
10 "	1,333	389	29.2
15 "	1,414	408	28.9

Pollen treated as above was used to fertilise selected stigmas. Though pollen, treated with different dosages germinated without much difference due to dosage, it will be noticed from the following table that the longer exposures were considerably less effective at fertilisation.

Duration of exposure to X-rays.	No. of flowers emasculated and pollinated.	No. of seeds set.	Percentage of seed-setting.
5 minutes.	48	20	41.7
10 "	39	7	18.0
15 "	61	11	18.0

It is remarkable that in the progeny raised from these seeds, one plant exhibited a suppression of the top eight leaf-blades showing only a stem merely ensheathed.

Millets Breeding Station, }
Coimbatore, 21-11-'34. }

G. N. Rangaswami Ayyangar.
V. Panduranga Rao.

ABSTRACTS

Report of the rice production and trade in the Madras Presidency By C. R. Srinivasan. The rice area and rice production have been generally on the increase during the last decade in almost all the rice producing countries of the east, particularly in Burma, Siam and Indo-China who on account of their huge exportable surplus, figure prominently in the international rice trade. Among the provinces of India, Madras occupies the fourth place as regards rice acreage and the third place as regards total production. The area under rice in Madras is about 11.4 million acres which form about 30 per cent of the area under cultivation. Of the 11.4 million acres, 8.2 millions are under irrigation while the balance is entirely dependent on rain. The net quantity of 5,036,322 tons of clean rice and 3,638,833 tons of millets produced annually in the province allows a consumption of 5 cwt. of cereals per year per adult head of the population which has been estimated to be 75 per cent of the total population. This quantity should be almost enough for the requirements of the Province, but since the yields of millets and the rice in the unirrigated areas are entirely dependent on favourable monsoons, there are big fluctuations in the production from year to year. There is usually a deficit of two to three hundred thousand tons of rice which she has to import from other countries and Provinces. Though this deficit was being met almost entirely by Burma previously, recently two other countries, Siam and Indo-China, have come into the field. The imports into Madras from these countries are continuously on the increase, and the import from Siam in particular is getting abnormally high since the beginning of 1934.

The entire import of Siam consists of broken rice, a bye-product in the preparation of raw rice for the international markets and is therefore very cheap. This cheapness combined with low transport charges by steamers is an inducement for the importers to bring it into the province in large quantities. Although its consumption is mainly confined to the poorer classes its price makes the price of the local produce come down relatively. There is no doubt that apart from the general economic depression gripping practically every country of the world, it is this huge import of the foreign rice into Madras that is responsible for disturbing the price levels of the local product. It so happens that the imports arrive at ports situated in the tracts which used to be the chief markets for the surplus product of the deltaic tracts in the province. In addition this imported rice is even sent to the centres in the interior up to distances of 80 to 100 miles from the ports where it competes with local rice. The rice or fall in price of the local rice follows more or less that of the imported rice. The price of rice in Madras at present is almost the same as that during the period 1904-08, i.e., it has come down by over 50 per cent since the depression began.

The railways are the chief means of transport of the local rice from place to place and the railway freight rates which had increased from 30 to 80 per cent during the last 15 years continue to-day at that level with certain exceptions, in spite of the fact that the value of the produce has come down considerably. Though the railway companies have allowed concession rates for the internal movement of the local produce wherever they come in competition with other means of transport, the advantages of such concessions cannot be fully availed of on account of the arrival of the cheap foreign rice at the ports and its distribution in the neighbourhood. Examination of the various agencies employed and the charges paid from the time the local produce is bought, up to the time it is offered to the consumer as a finished product gives no scope for a reduction in any of the items except the railway freights. Because of the high railway freights the movement of the produce within the province is getting more and more restricted and the foreign rice is gradually replacing it with the result that the condition of the rice grower is getting to be increasingly serious. The cost of production of rice was only 50 per cent of the market value of the produce before 1930, but now it has gone up to 70 to 80 per cent of the value, leaving the cultivator a much smaller margin to meet his other expenses from.

Besides the huge foreign imports, there is a certain amount of rice imported into the province from Bombay, Bengal, Central Provinces and it is not of much concern except probably the import from Central Provinces into the Vizagapatam district.

As regards foreign export from Madras, it amounted to only 75,504 tons in 1933-34 or roughly 1.5 per cent of the production. The exports are for Ceylon, Federated Malaya States and Strait Settlements. While the export from Godavari to Ceylon has considerably declined, that from Tanjore is being maintained and there is even a slight improvement since last year after the reduction in railway freights. The exports from Tanjore to Federated Malaya States and Strait Settlements are, however, again on the decline. The inland supply of rice from Madras, to the total extent of 2,402 (thousand maunds) in 1933-34 is for the markets of Mysore, Hyderabad and Bombay Deccan. This supply fluctuates with the production in these places. The long distance and the high transport charge by railways act as a deterrent against a larger movement than at present.

Examining the inland movement of rice within the province, the areas chiefly concerned in it are the districts of Godavari, Kistna, Nellore and Tanjore. The despatches from Northern Circars and Nellore are mainly in the form of raw rice though East and West Godavari districts send out some parboiled rice as well. In the Southern and Central districts, the movement is mainly confined to parboiled rice and paddy.

Machine milling of rice has made the hand pounding method obsolete in the rural areas. While rice is consumed as raw rice throughout the Circars and Ceded districts, in the South, Central and West Coast districts, it is mainly consumed as parboiled rice. The method of preparing parboiled rice varies according to the tracts and according to the size of the plant used in the manufacture. The movement of grain as paddy is confined mostly to places where the machine made parboiled rice is not popular. There is no doubt that it is the railway freight that stands in the way of greater internal movement of rice. Wherever the railway companies allowed concession rates the traffic has definitely increased.

Rice millers do not generally adopt any grade or standard for the produce they deal in. They mill rice according to the requirements of each market. A certain amount of adulteration between old and new, and superior and inferior rice, is always practised, particularly in the trade of raw rice. Good polishing

and the elimination of broken rice are practised only in the case of milling fine rice as it does not pay to adopt them in the case of coarse rice. In the trade there is no common unit followed. The produce is dealt, both in measure and in weight and the units of measure and weight vary from place to place often leading to confusion. The merit of the Departmental strains is appreciated by millers and merchants who pay a premium of two annas per bag of paddy for the produce obtained from them. It will be necessary to maintain their purity by the supply of pure stocks of seed from seed farms run in the different areas under Departmental supervision.

While the big landlords may be able to store their produce for some time after harvest, releasing it when price improves and even borrow money on the security of such stored produce, the large number of small cultivators have no such facilities, as they are obliged to sell off their produce immediately after harvest when price is low.

The following are some of the directions in which improvements may be introduced:—

- (i) Prohibition of the import of foreign rice and paddy or the imposition of a duty on such imports.
- (ii) Reducing the railway freights to the minimum for transporting rice and paddy from all the producing centres in the Province to the different consuming centres and retaining the maximum freight rates for transport of foreign imports arriving in the ports to inland centres,
- (iii) Forming associations of millers and merchants and using such associations to fix grades of produce, to control adulteration, and to introduce the system of purchase and sale by a standard unit of weight,
- (iv) Publishing regularly in the vernacular dailies of the Province, the market rates for rice and paddy at all the important centres of the Province,
- (v) Organization of co-operative purchase and sale societies with warehousing arrangements to help the small cultivators, and
- (vi) Hastening the spread of improved strains in all the rice tracts and intensifying the propaganda about better cultivation and manuring of the fields, so that the acre yields will be improved and the cost of production proportionately reduced. *(Author's summary).*

The Relation of Air Conditions to Tobacco curing. By Johnson and Ogden. (*Wisconsin Agri. Exp. Stn. Research Bulletin 110, July 1931*). By means of three constant air control chambers, operated simultaneously, the influence of various temperature and humidity conditions was determined on the rate of curing of tobacco and the quality of the product.

On this basis, the optimum temperature for curing was found to lie between 90 and 95°F., provided that a sufficiently high humidity was maintained at the same time. The ordinary range of constant temperatures which will permit of approximately normal curing, assuming favourable humidities at the same time, however, lies between 65° and 100°F.

The optimum relative humidity for curing is dependent upon the temperature. In general, at temperatures between 75° and 100°F., a percentage relative humidity corresponding to the temperature figure may be considered approximately satisfactory i.e., at 75, 80, 85, 90 and 95 per cent respectively.

Relative humidity is not as satisfactory a measure of the air-moisture relations surrounding curing tobacco as is the evaporating power of the air as measured by Livingston atmometers. On the latter basis, an evaporation of 8 to 12 cc. per day appears to be most desirable for all temperatures.

Detailed records of temperature and humidity conditions prevailing in the curing sheds have been taken over a period of six seasons. The average daily temperature over weekly periods rarely exceed 75°F., even at the top of the shed, indicating that ideal curing conditions are only for short periods of time. The average daily relative humidity for weekly periods was found to vary from as low as 44 per cent to as high as 96 per cent. The average daily evaporating power of the air for weekly periods has varied from about 3 cc. to 21 cc., a ratio of 1 to 7, indicating a much greater variation in actual air moisture relations than may be suspected from the records of relative humidity. Significant differences in temperature and humidity exist between the upper and lower tiers of the curing shed.

A preliminary report is made on the application of artificial heat to the air curing process, by means of a hot-air-furnace system installed in a curing shed. The practicability of the method will depend upon a number of factors yet to be determined.

Studies on the nature of "Pole-rot" and other forms of damage to the leaf occurring during the curing process have been aided by the use of toxic dusts such as the organic mercury compounds. The evidence from these studies points conclusively to the direct relation between a species *Alternaria* and "pole-rot".

(*Author's summary*).

Gleanings.

"Back to the Land"—Mr. Lloyd George's Precept and Example. When Indian Leaders advise Indian young men to go "back to the land," they do not set them an example by themselves becoming cultivators. But when Mr. Lloyd George spoke on the national importance of agriculture he meant business. Recently he and 100 Surrey farmers had been comparing notes on potatoes at a demonstration plot on his farm. He was overjoyed when it was found that his plot had come out best in the demonstration. He is more proud of his orchards than of anything else he has ever done. His eyes were bright as a child's when he plunged in among his heavily laden apple trees, to point out those that won him the first prize at the Crystal Palace, and others that had yielded fruit of unsuspected flavour and rich colouring. He went unerringly, obviously knowing them by heart, to other trees in the orchards that had so far been disappointments. "Remember those Worcesters you saw here last time?" he said; "No good at all. I was thinking of giving them up. Well, this year they have done wonderfully. Won the first prize." The farm at Churt is not the mere plaything of an elder statesman. It is a definite contribution to agriculture. For it was derelict when he took it, abandoned on account of its low grade soil, third or fourth class. Mr. Lloyd George is showing that it can yield abundantly when properly used. Mr. Lloyd George is cultivating about 240 out of 500 acres. Before his time the farm employed three men. To-day it employs thirty-one, earning from 45 s. to 50 s. per week. "I am happy," he said, "happier than I have ever been in my life. Politics were fun. I enjoyed all that; but they weren't somehow as satisfying as this is. I am addressing you not as an ex-Prime Minister. but as a registered potato-grower. I have ceased to be a politician. I have now become a cultivator of the soil; so that what I say is without reference to politics, and merely my observations as a cultivator of land. "I am opposed to the policy of restricting production merely to raise prices. It suits me personally, but in the long run it won't do any good." (*The Modern Review* for November, 1934, page 634.)

World Search for Pasture-grasses. Drought-resistant grass species, to be used in rebuilding the depleted rangelands of the west, are to be sought in

Central Asia by an expedition being sent out by the U. S. Department of Agriculture. On the edge of the Gobi Desert there are great natural grasslands, which have been pastured for thousands of years by nomad tribes, without any sign of exhaustion. In this region the temperature ranges from 100 degrees Fahrenheit in the summer to 40 degrees below zero in the winter, and severe droughts are frequent. Yet the grasses survive, and the herds of livestock and game thrive on them. The leader of the expedition will be Professor Nicholas Roerich, veteran explorer of interior Asia, from Kashmir to Altai mountains. With him will be his son, George Roerich, an expert in Central Asiatic languages, and two U. S. Department of Agricultural specialists in grasses H. G. MacMillan and J. L. Stephens. (*Science* Vol. 80. No. 2068. Supp. p. 8).

Wheel and Bearing Equipment for Farm Wagons and Trailers. Studies of the draft of wheel equipment for farm wagons and trailers conducted at the Ohio Experiment Station are briefly summarized. The wheels were tested on meadow, cultivated soil, cinders, gravel, and concrete roads. The net loads varied from 2,000 to 5,000 lb., and the rates of travel from 25 to 20 miles per hour. The wheel tires were of steel and rubber.

The cultivated soil required the greatest draft, followed in order by meadow, gravel, cinder, and concrete roads. On four of the five tractive surfaces and at all loads and rates of travel, the low pressure pneumatic tyre required less draft than any of the other types of wheels tested. The only exception was on concrete road, where there was very little difference between the steel wheel and the low-pressure pneumatic tyre. The greatest difference in draft between the low-pressure pneumatic tyre and the steel wheel occurred on gravel roads.

The importance of wheel diameter depends largely upon the tractive surfaces over which the load is transported. In cultivated soil, however, the smaller diameter wheels required more draft, ranging from 1 to 15 per cent more and depending upon the weight of the load transported and the rate of travel. On hard-surfaced roads or even in meadow, very little difference in draft was noted.

The width of rim seems to be a factor in cultivated soil only. With loads weighing from 5,000 to 6,000 lb. the 6 in. rim gave a slight advantage. The 5 in. rim excelled at all loads weighing below 5,000 lb. Wheels equipped with roller bearings show a decided decrease in the draft of the wagon. On cinder road the wagon skein type of bearing required the greatest draft, while the taper-roller bearing required the least. (*Experiment Station Record* Vol. 71, No. 2, Aug. 1934, Page 255.)

Development of Root System of Tobacco under varied conditions, with particular reference to Cultural Practice and Fire-holding Capacity. Roots of tobacco were examined at several stages in their development by excavating trenches at various distances and removing the soil in successive layers, 10 cm. wide and 20 cm. deep. Washing the soil from the roots was also employed. In transplanting the 6 wk-old seedlings from the nursery beds the tap root is usually broken so that nearly all the roots arising later belong to the secondary root system. A close relationship exists between growth of new roots and soil moisture. Transplanting is usually done at the end of the dry monsoon. Thus watering for a period of 10 days is necessary for root establishment. Although growth of these roots may continue slowly thereafter, yet no new roots arise until the plants are again irrigated or rains occur. Both rate of growth and the degree of branching were greater in wet than in dry soil (no field soil water-content determinations are given). The number of roots and the depth of penetration were also greater in wet and irrigated soils than in soil unirrigated and unwatered by rain. The dry, heavy, clay soil consists of hard clods into which the roots cannot penetrate; consequently, they develop chiefly in the crevices

between the clods, and are distorted, thick woody and sparingly branched. Upon the advent of rains the colloidal soil swells and largely or entirely closes the crevices, cutting off the air supply to these deeper roots, which die, probably as a result of oxygen deficiency. In the meantime a new growth of fine, densely branched roots has spread horizontally in the surface soil. Not enough water is applied in irrigation to close the crevices and pores of the clay soil. In sandy soil irrigation or rains cause an increase in the number of roots both in the shallower and deeper soil, nor is there normal oblique direction of growth appreciably changed. Roots served by tillage operations branch freely in wet soil but not at all in dry, where they often wither and die. Manuring with sulphate of ammonia and green manure (*Crotalaria anagyroides*) resulted in a greater number of adventitious roots, their more rapid growth and more profuse branching. The direction of growth was scarcely affected and the depth of penetration only slightly increased. The roots of banana (*Musa*) spread widely just beneath the soil-surface and none penetrated beyond the first foot. Applications of phosphoric acid had no effect upon depth of root penetration. Greater drought resistance of tobacco strain (EK) over strain (Y-10) was shown to be due to more adventitious roots, more extensive root branching, a higher rate of root growth, and a more thorough occupation of the soil. Maximum depths of penetration were 84 and 70 cm respectively. Root habits in regard to cultural practice are discussed. Irrigating soils of high chloride content in the surface soil will stimulate the absorption of too much chloride and in consequence will reduce the fire-holding capacity of the tobacco. Hence in such soils root development must be checked until rains set in, which dilute or leach the chlorides. Since irrigation does not bring these results about, or, if so, only temporarily, even fertilizers free from chlorides may harm the fire-holding capacity by stimulating the development of the absorbing area of the root system during the dry season. (*Biological Abstracts* Vol. 8 (5)—No. 11723—1934).

Crop and Trade Reports.

Paddy-Crop, 1934-35—Second forecast report. The average of the areas under paddy in the Madras Presidency during the five years ending 1932-33 has represented 13.3 per cent. of the total area under paddy in India.

The area sown with paddy up to the 25th November 1934 is estimated at 9,711,000 acres. When compared with the area of 10,491,000 acres estimated for the corresponding period of the previous year, it reveals a decrease of 7.4 per cent. The decrease in area occurs in all districts outside the Circars (Vizagapatam excepted) and Kurnool and is due to the receipt of late and insufficient rains. The first crop has been generally harvested throughout the Presidency. Normal yields have been reported from all districts except Guntur, Bellary, Anantapur, Cuddapah, Nellore, Chingleput, Salem, Coimbatore, Trichinopoly, Tinnevely and Malabar. The seasonal factor for the Presidency works out at 98 per cent. of the average as against 97 per cent. in the corresponding period of the previous year. The wholesale price of paddy per imperial maund of 82-2/7 lb. as reported from important markets towards the close of November 1934, was Rs. 2-14-0 in Madura, Rs. 2-13-0 in Nandyal and Vellore, Rs. 2-11-0 in Nellore and Salem, Rs. 2-10-0 in Tinnevely, Rs. 1-14-0 in Rajahmundry, Rs. 1-13-0 in Berhampur, Rs. 1-11-0 in Cocanada, and ranged from Rs. 2-0-0 to 2-6-0 in the other markets. When compared with the prices reported for October 1934, these prices are stationary in Berhampur, Vizianagaram, Nandyal, Vellore, Nellore, Tinnevely and Cochin; they have risen by 4 to 5 per cent. in Vizagapatam, Salem and Kumbakonam and are lower by 28 per cent. in Cocanada, 3 per cent. in Madura and 5 to 20 per cent. in the other markets.

Groundnut crop, Madras—1934—Intermediate report. The condition of the winter crop of groundnut is satisfactory in Vizagapatam, Chingleput, South Arcot, Chittoor and North Arcot. The crop has been affected by late sowing and subsequent heavy rains in Ganjam and by drought to some extent in the other districts, especially in the light soils of Bellary where it is reported to have suffered very much.

The wholesale price of groundnut (shelled) per imperial maund of 82-2/7 lb. as reported from important markets towards the close of October 1934 was Rs. 4-8-0 in Cuddalore, Rs. 4-5-0 in Vizagapatam and Vizianagaram and ranged from Rs. 3-8-0 to Rs. 3-13-0 in the other stations. When compared with the prices of the previous month, these prices reveal a rise of 19 per cent in Vizianagaram, 3 per cent in Vizagapatam and a fall of 13 per cent in Nandyal, 9 per cent in Cuddapah, 8 per cent in Cuddalore and 5 per cent in Salem.

Gingelly Crop—1934-35—Intermediate report. The yield of gingelly is expected to be normal only in Madura, Ramnad, Tinnevely and the West Coast. The crop has been affected to some extent by drought in most of the other districts.

Sugarcane Crop—Madras—1934. Intermediate report. The sugarcane crop was affected to some extent by drought in Bellary, Anantapur and Salem. The rains received towards the end of October and the beginning of November have lodged the crop in parts of the Cuddapah taluk. The condition of the crop is satisfactory in the other districts and the yield is expected to be normal if the season continues to be favourable.

The wholesale price of jaggery per Imperial maund of 82-2/7 lb. as reported from important markets towards the close of November 1934 was Rs. 10-5-0 in Erode, Rs. 7-2-0 in Trichinopoly, Rs. 6-5-0 in Nandyal, Rs. 6-3-0 in Bellary, Rs. 6-2-0 in Rajahmundry, Rs. 5-13-0 in Ellore and Cuddapah, Rs. 5-8-0 in Vizagapatam and Rs. 4-14-0 in Guntur and Vellore. When compared with the prices of the previous month, these prices reveal a rise of 2 per cent in Trichinopoly and a fall of 34 per cent in Guntur, 32 per cent in Vellore, 24 per cent in Nandyal, 21 per cent in Cuddapah, 19 per cent in Ellore, 16 per cent in Bellary, 11 per cent in Rajahmundry and 3 per cent in Vizagapatam. The price remained stationary in Erode.

College News & Notes.

Students' Corner. With the terminal examinations in progress, athletic activities were not very much in evidence during the month, on the college *maidan*; The Students' Club however participated in the Inter-District Olympic Sports held in Coimbatore and had the credit of knocking off the big cup—The Rao Bahadur Thiruvengadaswamy Mudaliar Cup—for the highest aggregate in the open events. They also entered for the Health Week Tournament in Foot ball and ran out winners. Of local interest, the final match in the Krishnamoorthy Rao Memorial Inter-Tutorial Hockey Tournament resulted in a win for Mr. P. V. Ramiah's wards who met and beat Mr. Sundararaman's.

The College Day Celebrations. The College Day Celebrations came off on the 13th (Conference) 14th (Business meeting and Entertainments) and 15th (Sports) of December. A detailed description will be published in the next issue.

Farewell. Rao Sahib V. Muthuswamy Ayyar, Lecturer in Agriculture, who retired from active service on 16th December was the recipient of a number of 'teas' arranged in his honour, by the students, the officers' club, the Indian Officers' Association and by other friends. Mr. K. Raghavacharya who was posted as lecturer in Agriculture, in his place assumed charge of his duties from 16th.

Visitors. Mr. R. C. Burt, Acting Vice-chairman of the Imperial Council of Agricultural Research, Mr. Hutchinson, Plant Geneticist, Indore, Mr. P. H. Rama Reddy, Secretary, Indian Central Cotton Committee, and Mr. S. V. Ramamurthy, Director of Agriculture, were amongst those who visited the College during the month, in addition to those who attended the College Day and Conference.

Association of Economic Biologists. Under the auspices of the above Association, Mr. Hutchinson delivered a very interesting lecture on the 9th on the "Importance of Genetics to the Crop Breeder."

Weather Review (NOVEMBER—1934)

RAINFALL DATA

Division	Station	Actual for month	Departure from normal	Total since January 1st	Division	Station	Actual for month	Departure from normal	Total since January 1st	
Circars	Gopalpore	2.7	-1.3	4.7	South	Negapatam	8.9	-8.8	34.9	
	Berhampore *	0.8	-6.1	33.2		Aduthurai *	4.8	-5.6	30.4	
	Calingapatam	0.4	-3.5	25.0		Madura	2.9	-2.1	24.7	
	Vizagapatam	4.1	+0.3	33.1		Pambar	4.2	-7.8	30.1	
	Anakapalli *	3.6	-1.7	34.7		Koilpatti *	0.7	-6.3	23.1	
	Samalkota *	2.3	-1.3	36.0		Palamkottah	0.6	-6.8	23.9	
	Maruteru *	1.4	-3.7	28.4		West Coast	Trivandrum	6.5	-0.1	56.2
	Cocanada	4.1	-1.3	40.3			Cochin	3.1	-3.5	97.9
	Masulipatam	7.7	+2.0	35.1			Calicut	6.2	+0.9	107.9
	Guntur *	1.7	-1.9	33.5			Pattambi *	4.3	+0.5	89.7
Ceded Dists.	Kurnool	2.0	+0.9	20.3	Taliparamba *		4.4	0.0	131.0	
	Nandyal *	1.5	0.0	27.0	Kasargode *		3.0	-1.0	121.4	
	Hagari *	1.6	-0.2	14.8	Nileshwar *		3.2	-0.2	126.5	
	Bellary	1.0	-1.2	14.4	Mangalore		1.4	-1.7	116.1	
	Anantapur	0.3	...	9.2	Mysore and Coorg		Chitaldrug	0.6	-1.6	26.8
	Cuddapah	3.2	-0.4	23.3			Bangalore	1.1	-1.9	24.5
Carnatic	Nellore	2.4	-8.9	19.9		Mysore	3.7	+1.2	36.3	
	Madras	1.9	-12.4	34.5		Mercara	2.0	-1.2	109.4	
	Palur *	7.9	-3.4	53.9		Hills.	Kodaikanal	5.1	-3.1	53.9
	Palakuppam *	2.8	-8.1	35.6			Coonoor	5.4	...	51.1
Cuddalore	10.2	-4.9	52.2	Ootacamund *	4.7		-0.3	43.7		
Central	Vellore	7.2	+0.3	40.4	Nanjanad *		2.5	-1.5	40.0	
	Hosur cattle farm *	2.3	0.8	24.7	Summary of General Weather Conditions.					
	Sulem	3.6	-0.2	34.4						
	Coimbatore	1.1	-2.7	19.6						
	Coimbatore Res. Inst. *	2.7	-1.8	22.2						
Trichinopoly	6.1	+0.5	25.7							

* Meteorological Stations of the Agricultural Department.

Summary of General Weather Conditions.

Weather remained dry over the major portion of the country after the 10th. Three depressions appeared during the month causing generally local rain in different portions of the country. The unsettled conditions off Konkan which were forming towards the end of last month developed into a storm on the 1st centred within one degree of Lat. $14\frac{1}{2}^{\circ}$ N and Long. $71\frac{1}{2}^{\circ}$ E. This caused widespread rain in Konkan and Deccan and local rain in Malabar, North Madras and Central Provinces. The storm moved northwards on the 4th and got filled up on

the 5th off Kathiwar. A second depression appeared on the 5th in the South East Bay of Bengal and moving North-Northwest developed into a storm and crossed the Arakan coast near Akyab on the 10th November. It weakened after causing widespread rain in South East Madras, Chota Nagpur, Bihar, Bengal and Assam. Locally heavy rain occurred in Burma and the Chittagong coast. After the 10th weather remained generally dry over the country except for a few falls in Tennesserium and South Madras. The third depression appeared in the Bay on the 25th and developed into a storm centred within a degree of Lat. 10°N and Long 84°E on the 26th, causing local rain in South East Madras, the storm crossed the coast near Negapatam on the 28th, weakened and became unimportant on the 29th. This caused local rain in South East Madras, Mysore and a few falls along the North Madras coast.

Four western disturbances affected the extreme north on the 6th, 9th, 16th and 20th causing a few showers in Kashmir and Baluchistan and passed away eastwards.

Rainfall was markedly below normal in the whole of the Presidency, that in Madras being in defect by—12·4." Night temperatures were below normal over the Peninsula during the dry weather period.

Weather Report for the Research Institute Observatory :

Report No. 11/34.

Absolute maximum in shade	88·5° F
<i>Absolute Minimum in shade</i>	57·5° F
Mean maximum in shade	84·6° F
Departure from normal	- 0·1° F
Mean minimum in shade	66·5° F
Departure from normal	- 2·2° F
Total rainfall	2·74"
Departure from normal	- 1·79"
Heaviest fall in 24 hours	1·19"
Total number of rainy days	4
Mean daily wind velocity	3·3 M. P. H.
Mean humidity at 8 hours	74·2%
Departure from normal	- 7·1%
Total hours of Bright Sunshine	202·1
Mean daily hours of Bright Sunshine	6·7

General Summary. Rainfall is in defect by -1·79". Minimum temperature is below normal. Humidity is in large defect.

ADDITIONS TO THE LIBRARY, OCTOBER 1933

A. Books.

1. Forage Plants and their Culture. *Piper C. V.* (1931).
2. The Profitable Culture of Vegetables for Market Gardens, Small Holders and others. *Thos. Smith* (1932).
3. The Culture of Vegetables and Flowers from seeds and Roots. 18th Edn. *Sutton & Sons* (1930).
4. The New Illustrated Gardening Encyclopaedia. *Sudell R.* (1932).
5. Methods and Materials of Chinese Table Plant Culture. *McClure F. A.* (1933).
6. Spices and Condiments. *Redgrove M. S.* (1933).
7. Roadside Planting. *Road Beautifying Association Pub.* (1930).
8. World Weather—Part V—Tabulation. *Bliss E. W.* (1932).
9. Handbook of the Collections Illustrating Pumping Machinery—Part I—Historical. *Westcott G. F.* (1932).
10. England Agricultural Marketing Act of 1933. (1933).
11. Dairy Cattle: Selection Feeding and Management. *Yapp W. W. & Nevens W. B.* (1930).
12. Geese: Breeding Rearing and General Management. *Appleyard R.* (1932).
13. Practical Sheep Farming. *Norris T. C.* (1933).
14. Sex Determination. *Crew F. A. E.* (1933).
15. Studies on the Development and Situation of Pedigree Registering in the Cattle Breeding Industry. *Engeler W.* (1931).
16. A Text Book of the Diseases of the Small Domestic Animals—2nd Edn. Revised. *Brumely O. V.* (1931).
17. Vitamins and their Dietary Essentials. *Aykroyd W. R.* (1933).
18. The Relation of Food to Diseases. *Dixon S.* (1932).
19. Cytological Technique. *Baker J. R.* (1933).
20. Plant Ecology. *Leach W.* (1933).
21. Gramineae—Setaria (Flora of Tropical Africa—Vol. IX—Pt. 5). *Prain D.* (1930).
22. The Green Leaf: The Major Activities of Plants in Sunlight. *Mac Dougal D. T.* (1930).
23. Flowerless Plants—Part II of an Introduction to Structural Botany—10th Edn. Revised. *Scott D. H.* (1932).
24. Proceedings of the Sixth International Congress of Genetics Vols. I & II. (1932).
25. An Introduction to Biology. *Cole E. C.* (1933).
26. The Action of the Living Cell: Experimental Researches in Biology. *Turck F. B.* (1933).
27. Evolution, Fact and Theory (Compilation). *Wells H. G. Huxley J. S. & Wells G. P.* (1932).
28. Fighting the Insects: The Story of an Entomologist. *Howard L. O.* (1933).
29. Insects: Man's Chief Competitors. *Flint W. P. & Metcalf C. L.* (1932).
30. Recent Advances in Pathology. *Hadfield G. & Garrod L. P.* (1932).
31. Phytopathological and Botanical Research Methods. *Rawlins T. E.* (1933).
32. A Manual of Agricultural Chemistry—5th Ed. Revd. *Ingle H.* (1933).
33. Chemical Encyclopaedia: A Digest of Chemistry and its Industrial Applications—5th Edn. *Kingzett C. T.* (1932).
34. Kostychev's Chemical Plant Physiology. *Lyon C. J. Tr.* (1931).
35. Low temperature Carbonisation: Narrow Brick Retorts at the Fuel Research Station. (1933).
36. An Essay on the Nature and Significance of Economic Science. *Robbins L.* (1932).
37. Muscular Work, Fatigue and Recovery. *Crowden C. P.* (1932).
38. Pitman's Commercial Atlas of the World. *Rutter W. P.* (1932).
39. The Common Sense of Political Economy and Selected Papers and Reviews of Economics. *Wicksteed P. H.* (1933).
40. The Social Economics of Agriculture. *Gee W.* (1932).
41. Uses of Waste Materials: The Collection of Waste Materials and their Uses for Human and Animal Food, in Fertilizers and in Certain Industries 1914-1922. *Bruttini A. B.* (1923).
42. Soviet Economics (Trans. by M. Campbell). *Ed. Dobbert G.* (1933).
43. Reconstruction and Education in Rural India. *Premchand Lal.* (1932).
44. What is Probable Error. *Tocher J. F.* (1931).
45. Tables for Statisticians and Biometricians, Vol. I and II. *Pearson K. Ed.* (1931).
46. The Indexing of Books and Periodicals. *Walsh J. W. T.* (1930).
47. Readings in Library Methods. *Arnett L. D. & Arnett E. T.* (1931).
48. Basic German for Science Students with Vocabulary and English Translations of the German Passages. *Barker M. L.* (1933).
49. Madras District Gazetteer: Anantapur District.
50. Cuddapah District.
51. Chingleput District. Vol. III. (1933).

B. Reports.

1. Detailed Administration Reports of the Agricultural Chemist, Mycologist and Entomologist of the Department of Agriculture, Madras for 1932-33, 1(a). Administration Report of the Department of Agriculture and Fisheries, Travancore, for the year 1931-32. 2. Annual Report of the Indian Central Cotton Committee, Bombay for the year ending 31st August 1932. 3. Report of the Rothamsted Experimental Station for 1932. 4. Fourteenth Annual Report—Agricultural Census 1933 of the Colony and Protectorate of Kenya. 5. Report on the Agricultural Department Antigua for 1932. 6. Report on Higher Education in the State of New York for the School Year Ending July 31, 1931. 7. Forty-Second Annual Report of the Storrs Agricultural Experiment Station (Connecticut) for the year ending June 30, 1930. 8. Forty-Third Annual Report of the Storrs Agricultural Experiment Station (Connecticut) for the year ending June 30, 1931. 9. Annual Report of the Maine State Pomological Society for 1931. 10. Report of the Hawaii Agricultural Experiment Station for 1932. 11. Summary Report of Progress of the Maine Agricultural Experiment Station for 1932. 12. Abstract Proceedings of the Twenty-Sixth Meeting of the Indian Central Cotton Committee, Bombay held on the 2nd and 3rd February 1933. 13. The Hokkaido Imperial University Calendar for 1933-1934.

C. Bulletins, Memoirs &c.

14. Tamil Sayings and Proverbs on Agriculture. *Mad. Agri. Dept Bull.* 29, 1933 New Series. 15. Spraying of Coffee in South India. 16. Annual Report of the Coffee Scientific Officer, 1932-33. *Mysore Coffee Expt. Stn. Bull.* 9, 10. 17. Flora Arabica—The Botanical Exploration of Arabia. *Records of Bot. Surv. of Indian Vol. VIII—No. 5.* 18. Registration of Earth Current with Natural Electrodes. *India Meteorol. Dept. Mem. Vol. XXVI, Part 1.* 19. Coir—Report on the Attributes and Preparation of Coconut Fibre. 20. Banana Storage—An Account of Recent Investigations into the Storage Behaviour of Several Varieties. *Imp. Mark. Bd. Pub. Nos. 71, 72.* 21. West Cambridgeshire Fruit Growing Area. *Eng. Mini. Agri. & Fish. Bull.* 61. 22. Some Cultivated St. Lucia Soils. *Eng. Imp. Bur. Soil. Sci. Tech. Comm. No. 4.* 23. The Determination of the Inorganic Colloidal Constituents of Soils. 24. Winogradsky's Soil Plaque Method for Determining the Nutrient Requirement of Soils. 25. Pollen Analysis. 26. Work at Rothamsted Experiment Station. 27. The Problem of the Pakihi lands of New Zealand. 28. The Measurement of Soil Colour. 29. Soil Research at the Imperial College of Tropical Agriculture, Trinidad. 30. Soil Research at the Waite Institute Glen Osmond, South Australia. 31. Soil Research in Progress at Imperial Institute of Agricultural Research, Pusa, India. 32. Soil Research in Progress at the Imperial Research Institute of Ceylon. 33. Investigations relating to Soils and Fertilizers at the Cawthorn Institute, Nelson, New Zealand. 34. Reclamations of the Pontine Marshes. 35. Cajander's Theory of Forest Types. 36. Sub-irrigation. 37. The Work of the Chemical Section, Department of Agriculture, Nigeria. 38. A Novel Container for Soil Monoliths. 39. Irrigation and Engineering Problems in Sind. 40. (1) Saline and Alkali Soil Reclamation. (2) Sulpho cyanide as a Weed Killer. (Extract). *Eng. Imp. Bur. Soil. Sci. Monthly Letter Nos. 1, 2, 5, 6, 7, 8, 9, 11, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24.* 41. A List of Insects with their Parasites and Predators in Malaya. *S. S. & F. M. S. Agri. Dept. Sci. Ser. Pub. No. 13.* 42. Reports of the Research, Economic and Agricultural Education Branches for the year 1932. *S. S. & F. M. S. Agri. Dept. Genl. Ser. Pub. No. 14.* 43. Economic Investigation into Live-Stock Farming in the Northern Transvaal, 1927-30. *Union of S. Africa Agri. Dept. (Econ. Ser. No. 17) Bull. No. 121.* 44. The White Grubs or "Hardback Beetles" Injurious to Sugarcane in British Guiana. *British Guiana Agri. Dept. Ent. Bull. 2.* 45. Etiology of the Chocolate Spot

- Disease of Broad Beans. *New South Wales Agri. Dept. Sci. Bull.* 43. 46. Basic Law for the Agricultural Reform. *Spain Institute of Agricultural Reform Pub.* 47. Cotton Fibres. III Cell Divisions in the Epidermal Layer of the Ovule subsequent to Fertilization. *Boyce Thompson Inst. for Plant Research Separate Vol.* 5-2. 43. Properties and Uses of Calcium Cyanamide. *Boyce Thompson Inst. Professional Paper—Vol. 1—24.* 49. Rice Growing in the United States. A Preliminary Report. *Iowa State College Pub. April 1933.* 50. Tree Planting, Soil Blasting, Tree Rejuvenation, other Farm Uses. *Hercules Powder Co., Pub.* 51. Drying on the Farm. *Brit. Crop. Driers Ltd. Pub.* 52. A Contribution to the Knowledge of Regeneration in Higher Plants. *Hokkaido Imp. Univ. Jrl. Faculty of Agri. Vol. XXXV—Pt 1.* 53. The Control of Fungus Diseases. *Cyprus Agri. Dept. Bull. 1 (Mycol. Series).* 54. Picking, Grading and Packing of Citrus Fruits—A Guide for Exporters. *Cyprus Agri. Dept. Bull. No. 2 (Horti. Series).* 55. Range Grasses of Hawaii. 56. Survey of the Physical Features that affect the Agriculture of the Kona District of Hawaii. 57. Cane Molasses in Poultry Rations. *Hawaii A. E. S. Bull. Nos. 65, 66, 67.* 58. Experiments with Fertilizers on Coconut Palms and Variation in Palm Productivity *Porto Rico A. E. S. Bull. No. 34.* 59. Vitamin D in the Nutrition of the Dairy Calf. *Wisconsin A. E. S. Res. Bull. 115.* 60. Solar Water Heaters. *Florida A. E. S. Bull. 68.* 61. Cotton Root Development in Certain South Louisiana Soils. 62. Retail Prices of Fertilizer Materials Cornell and Mixed Fertilizers. *Louisiana A. E. S. Bull. Nos. 232, 545.* 63. Terracing to Control Erosion—A Profitable Farm Practice. *Kansas A. E. S. Extn. Bull. No. 70.* 64. The Influence of Various Systems of Grape Training on Fruit Production. 65. The Relative Value of Different Carriers of Phosphorus. 66. The Pectic Constituents of Tomatoes and their relation to the Canned Product. 67. Does Tile Drainage Pay? 68. The Effect of Sulphur on Soils and on Crop Yields. 69. Environmental Factors Influencing Wheat Production in Maryland. 70. Physiological Shrinkage of Potatoes in Storage. 71. Fertilizer Studies and the Production of Sweet Potatoes. 72. Organization and Management of Maryland Farms—Piedmont Plateau Region. 73. Economic Efficiency of the Farm Layout in Maryland. 74. Taxation in Maryland with Special Reference to Agriculture. *Maryland A. E. S. Bull. Nos. 250, 289, 291, 295, 296, 297, 303, 311, 337, 338, 339* 75. Agricultural Land Classification and Land Types of Michigan. 76. The Michigan Pear Industry: Its Status and Trends. 77. Experimental Studies in Feeding Fattening Lambs. 78. Spraying and Dusting Potatoes in Michigan. 79. Motor Truck Marketing of Michigan Livestock. *Michigan A. E. S. Spl. Bull. Nos. 231, 232, 233, 234, 235.* 80. A Study of the Ash Constituents of Apple Fruits During the Growing Season. 81. The Uses of Formaldehyde Dust in Growing Seedlings. 82. Roadside Marketing of Agricultural Products by Ohio Farmers. 83. Control of Alternaria Blight of Ginseng with Bordeaux Mixture and Injuries Accompanying its use. 84. Sources of Market Milk and Butter-fat in Ohio. 85. The Potato Scab-gnat, *Pnyxia Scabiei* (Hopkins). *Ohio. A. E. S. Bull. Nos. 519, 520, 521, 522, 523, 524.* 86. Viability of Eggs and Larvae of the Apple Maggot. 87. Bacteriological Studies of a High Temperature, Short-Time Pasturiser. 88. Notes on the Species of Apples 1. The American Crabapples. 89. Chromosome Numbers in Aneuploid Apple Seedlings. *New York A. E. S. Tech. Bull. Nos. 206, 207, 208, 209.* 90. New or Noteworthy Fruits. 91. Abnormal Enlargement of Peas From Plants affected with Root-rot 92. Some Carbohydrate Changes in Shelled Green Peas. 93. Experiments with Commercial Nitrogenous Fertilizers on Apple Orchards. 94. Evaluation of Applications of Lime-Sulfur for the Control of Apple Scab. 95. Raspberry Growing in New York State Cultural Practises and Disease Control. 96. The Laboratory Detection of Bovine Mastitis. 97. Codling Moth Control. 98. Sterilization of Ice Cream Freezers. *New York A. E. S. Bull. Nos. 620, 621, 622, 623, 624, 625, 626, 627, 628.* 99. Farming Systems for Eastern Washington and Northern Idaho. *Idaho A. E. S. Bull. No. 244.* 100. Lead and Arsenic

Spray Residue Removal from Apples. *Idaho A. E. S. Tech. Bull.* 286. 101. Effect of Manufacturing and Preserving Processes on the Vitamins of Cranberries. 102. The Effect of Certain Cropping Systems on the Yield and Quality of Havana Tobacco. 103. The Significance of Length of Biological Laying Year in Production Breeding. *Massachusetts A. E. S. Bull. Nos.* 296, 297, 293. 104. Studies in the Genetics and the Cytology of *Ustilago avenae* and *Ustilago levis*. *Minnesota A. E. S. Tech. Bull. No.* 87. 105. Relation of Variations in the Human Factor to Financial Returns in Farming. 106. Filling Silos with the Field Ensilage Harvester. 107. Crop Investigations on Sandy Lands. 108. Crop Investigations on Peat Lands. *Minnesota A. E. S. Bull. Nos.* 288, 290, 291, 292. 109. (1) Seed Transmission and Sterility Studies of Two Strains of Tobacco Ringspot. (2) A Virus Disease of Delphinium and Tobacco. (3) A Virus Disease of Plum and Peach. 110. The Resistance of Certain Red Clovers and Alfalfas to Leafhopper Injury. *Kentucky A. E. S. Res. Bull. Nos.* 327, 329. 111. Soils and Fertilizer Experiments, Experiment Station Farm, Lexington. *Kentucky A. E. S. Bull. No.* 331. 112. (1) Winter versus Summer Feeding of Corn to Heavy Steers Finished on Grass. (2) Wheat as a Fattening Feed for Yearling Steers. *Kentucky A. E. S. Bull. No.* 332. 113. Infections of Fetuses and Foals. 114. The Comparative Effect of Muriate and Sulphate of Potash on the Composition and Quality of White Burley Tobacco. *Kentucky A. E. S. Res. Bull. Nos.* 333, 334. 115. Changes in Quantity and Composition of Fat in Hogs Fed a Peanut Ration followed by a Corn Ration. 116. Effectiveness of the Swine Sanitation System in the South. 117. Character and Behaviour of Organic Soil Colloids. *U. S. Agri. Dept. Tech. Bull. Nos.* 368, 374, 377. 118. Farm Blumbing. 119. Strawberry Diseases. 120. Beef Production on the Farm. 121. Feeding Dairy Cows. 122. Milk for the Family. *U. S. A. Farm. Bull. Nos.* 1426, 1458, 1592, 1626, 1705.

D. Circulars, Leaflets &c.

123. Cotton Research in the Punjab. 124. Some Problems of Cotton Growing and Its rotation crops in the Central Provinces and Berar and the work hitherto done on them by the Botanical Section, Nagpur, and at the Akola Farm, under the auspices of the Indian Central Cotton Committee, Bombay. 125. The Improvement of Broach Cotton Research Work conducted at the Government Farm, Broach. *Ind. Cent. Cotton Com. Pub.* 127. Wire Fencing for Grassland. *Eng. Mini. Agri. & Fish. Adv. Leaf.* 186. 128. The Grading of Western Australian Timbers; Report on, and Suggested Specifications for, the Grading of Jarrah and Karri based on Investigations in 1932. 129. Investigations on the Buffalo Fly (*Lyperosia exigua* de Meij). *Australia Council for Sci. & Ind. Res. Pamphlet No.* 41, 43. 130. Design Factors Affecting the Strength and Rigidity of Wooden Crates. 131. Tests of Varieties and Strains of Large-seeded Virginia-Type Peanuts. 132. The Survival of European Corn-Borer Larvae in Barns and Other Storage Places. 133. Three Bacterial Spots of Tomato Fruit. 134. Grades of Peat and Muck for Soil Improvement. *U. S. Agri. Dept. Cir.* 236, 272, 281, 282, 290.

E. Reprints.

135. The World Rice Situation in June 1933. 136. Factors affecting Infection of Wheat Heads by *Gibberella Saubinetti*. 137. Sclerotia-Forming Habits of the Cotton Root-Rot Fungus in Texas Black-Land Soils. 138. Artificial Transmission of Sugarcane Mosaic. 139. Inheritance of *Fusarium* Resistance in Brussels Sprouts and Kohlrabi. 140. Effects of Chlorine, Bromine and Fluorine on the Tobacco Plant.

F. English Translations of Original Articles in Russian. (*Imperial Agricultural Bureau.*)

141. Komarov:—Origin of Cultivated Plants. (*Abr. Trans.*) 142. Hiltbrandt:—Full Summary of Sesame. 142. Hiltbrandt:—Geographical Distribution

of the Economical Characters of Sesame and the Utilization of its World Assortment under the conditions of U. S. S. R. 1933. 144. Hiltbrandt:— Full Summary of Sesame. (*Sesamum Indicum* L.)—another article. 145. Popova:— The Castor Bean in Central Asia. 146. Kappert:— On the Evaluation of Dihybrid-Mendelian Ratios in Linkage Studies.

NOVEMBER 1933

A. Books.

1. Fertilizers and Food Production on Arable and Grassland. *Keable F.* (1932).
2. Fruit cultivation in Ceylon and South India. *Taylor F.* (1933).
3. A Consolidated Record of the Field Experimental Works carried out by the Department of Agriculture in the Central Provinces between 1900-1930. *Allen R. G.* (1933).
4. The Genetics of Cattle—(Reprint from *Bibliographia Genetica*—Vol. X). (1933).
5. Studies in Determinate Evolution. (Reprint from *Ohio Journal of Science*. 1929-1933). *Schaffner J. H.* (1933).
6. The Evolution of the Class Insects. *Tillyard R. J.* (1930).
7. Handbook of Study and Travel for Indian Students going abroad. *Ranganadham P. D.* (1932)
8. The Layman's Law Guide. *Sequeira A. C.* (1932).
9. Madras District Gazetteers—Nilgiri District—Vol. III. *Madras Govt. Pub.* (1933).
10. Interest Calculation Tables and Ready Reckoners for use in the Account Offices. *India Govt. Pub.* (1933).

B. Reports.

1. Report of the Department of Industries, Madras for the year ending 31st March 1933.
2. Annual Report of the Royal Botanic Garden and the Gardens in Calcutta and of the Lloyd Botanic Garden, Darjeeling for the year 1932-33.
3. Report of the Botanical Survey of India for 1931-32.
4. Annual Report of the Kenya Colony and Protectorate—1932.
5. Report of the Minister of Agriculture of the Province of Ontario—1932.
6. Report of the Agricultural Department St. Kitts-Nevis—1932.
7. 15th Annual Report of the Federal Board for Vocational Education—1931.
8. 16th Annual Report of the Federal Board for Vocational Education for 1932.
9. Forty-Fourth Annual Report for the Fiscal year ending June 30, 1932 of the Arkansas Agricultural Experiment Station
10. Annual Report of Wisconsin Agricultural Experiment Station for 1931-1932 (Facts for Farmers).
11. Annual Report of the Extension Service—College of Agriculture—University of Wisconsin for 1932.
12. Annual Report of Delaware Agricultural Experiment Station for the Fiscal year ending June 30, 1932.
13. Progress Report of Alaska Agricultural Experiment Station for 1932.
14. Biennial Report of Utah Agricultural Experiment Station for July 1, 1926—June 30, 1928.
15. Biennial Report of Utah Agricultural Experiment Station for July 1, 1928—June 30, 1930.
16. Biennial Report of Utah Agricultural Experiment Station for July 1, 1930—June 30, 1932.

C. Special Publications.

17. Vegetables of New York: Peas of New York. *New York A. E. S. Pub. Vol. 1—Part 1.*
18. Vegetables of New York: Beans of New York. *New York A. E. S. Pub. Vol. I—Part. II.*
19. Bibliography of Tropical Agriculture 1932. *Rome International Inst. of Agri. Pub.* 1933.
20. Report of the Committee of Definitions: The Problem of Marketing Terminology. *Natms Bull. Pub. by National Asso. of Teachers of Marketing & Advertising.*
21. Ventilated Storage for Potatoes. *Penna Marble Lab. Pub.*
22. The Egyptian Agricultural Review: The Agricultural and Industrial Exhibition Special Number. *Egypt Mini. Agri. Pub.*
23. A Survey of Recent Work of the Department in relation to its bearing on Agricultural and Horticultural Practice.—March 1933. *Univ. of Bristol. Dept. of Agri. & Horti. Pub.*
24. The Progress in Forest Entomology and Protection against Forest Insects. *U. S. Agri. Dept. Senate Document No. 12—Separate No. 28.*

D. Bulletins, Memoirs &c.

25. Tamil Sayings and Proverbs on Agriculture. 26. Insects Injurious to Vegetable and Flower Plants in South India. *Madras Agri. Dept. Bull. Nos. 29 and 34 of 1933.* 27. The Foundations of Yarn--Strength and Yarn Extension. 28. A Note on the Early History of Cotton. 29. Ginning Percentage and Lint Index of Cotton in relation to the Number of Cotton Fibres per Seed, the Effect of Environment on Ginning Percentage and the Determination of Unit Fibre Weight. *Indian Cent. Cott. Comm. Tech. Ser. Nos. 7, 12, 13--Bull. Nos. 12, 17, 18.* 30. A Comparison of Some Methods of Testing the Breaking Strength of Single Cotton Fibres. 31. Studies in the Sampling of Cotton for the Determination of Fibre Properties. 32. The Weight per Inch of Fibres of Different lengths, and the Numbers of Fibres of Different Lengths per Seed, for each of the Standard Indian Cottons. 33. The Foundations of Yarn Strength and Yarn Extension III--The Clinging Power of Cotton. 34. Studies in the Sampling of Cotton for the Determination of Fibre Properties. 35. Fibre Length Irregularity in Cotton. *Indian Cent. Cott. Comm. Technological Bull. Ser. B. Nos 5, 6, 7, 8, 10, 16.* 36. Spinning Tests on Mixings of Long Staple and Short Staple Indian Cottons. 37. Technological Report on Banilla Cotton, 1930-32. 38. Variation in the Moisture Content on Baled Indian Cotton with Atmospheric Humidity. 39. Technological Reports on Standard Indian Cottons--1933. 40. Testing of Indian Cottons for Quality at the Technological Laboratory. *Indian Cent. Cott. Comm. Technological Bull. Ser. A. Nos. 20, 22, 23, 24, 25.* 41. A Further Note on the Colorimetric Estimation of Humic Matter in Mineral Soils. *Imp. Bur. of Soil Sci. Monthly Letter No. 45.* 42. Grazing: Papers read at the Meeting of the British Association for the Advancement of Science at Leicester on September 11th, 1933. 43. Technique Employed in Grassland Research in New Zealand. *Imp. Bur. Plant Genetics Herbage Plants Bull. Nos. 10, 11.* 44. Recent Researches on Helminth Immunity. *Imp. Bur. Agri. Paras. Pub. 1933.* 45. Studies on the Transport of Carbo-Hydrates in the Cotton Plant: III. The Polar Distribution of Sugar in the Foliage Leaf. *Trinidad Cotton Res. Stn. Mem. Ser. B. Physiology.* 46. Merino Wool Classing and Hints on Shed Work. 47. Fruit Production in the Union: The 1931-32 Deciduous Fruit Export Season. *Union of S. Africa Agri. Dept. Bull. Nos. 117, 122.* 48. Methods for Preserving Plants in their Natural Form and Colour. *Egypt Mini. Agri. Tech. & Sci. Bull. No. 124.* 49. Some Observations on the Citrus Industry of Palestine, with reference to Application of Improved Methods to the Citrus Industry of Cyprus. *Cyprus Agri. Dept. Bull. No. 1 (Horti. Ser.).* 50. Factors affecting the Price of Rice. 51. Pima Egyptian Cotton in Irrigated Rotations at the Yuma Field Station, Bard, Calif. 52. A Classification of North American Agallian Leaf Hoppers. 53. Microbiological Studies of Salt in Relation to the Reddening of Salted Hides. 54. Observations on the Thermal Death Points of *Anastrepha Ludens* (Loew). *U. S. Agri. Dept. Tech. Bull. Nos. 297, 369, 372, 383, 400.* 54(a). The Rough-Headed Corn Stalk Beetle in the Southern States and its Control. *U. S. Agri. Dept. Farmers' Bull. No. 875.* 55. The Composition of Soybean Plants at Various Growth Stages as related to their Rate of Decomposition and Uses as Green Manure. *Missouri A. E. S. Res. Bull. No. 173.* 56. Factors in the Organization and Successful Operation of Louisiana Rice Farms, 1930. *Louisiana A. E. S. Bull. No. 233.* 57. Farm Mortgage Foreclosures in Minnesota. 58. Diseases affecting Moose. *Minnesota A. E. S. Bull. Nos. 293, 294.* 59. Utilization of Agricultural Wastes and Surpluses. *Iowa A. E. S. Vol. XXXI--No. 43.* 60. Approved Practices for Peanut Growers. *North Carolina A. E. S. Bull. No. 281.* 61. A Study of Certain Economic Factors in relation to Social Life Among Oklahoma Cotton Farmers. *Oklahoma A. E. S. Bull. 211.* 62. Farm Lighting Systems. *Kansas State College Bull. Vol. XVI--No. 7.* 63. Sorghum Production in Kansas. *Kansas State College Bull. 265.* 64. The Effect of Fertilizers, Crop Rotation and Weather

Conditions on the Anchorage of Corn Plants. *Virginia A. E. S. Tech. Bull.* 46. 65. Reaction Studies of Delaware Soils. *Delaware A. E. S. Bull.* 155. 66. The Mechanism of Buffer Action in Soils. *Delaware A. E. S. Bull.* 166 & *Tech. No.* 11. 67. Insurance of Farm Families. *Michigan A. E. S. Tech. Bull.* 133. 68. Population Trends in Michigan. 69. Trends in Cherry Production in Michigan. *Michigan A. E. S. Spec. Bull. Nos.* 236, 237. 70. Muck Soil Management for Onion Production. *Michigan A. E. S. Extn.* 123. 71. The Influence of Nitrogen in Soil on Azofication. 72. Mutual Irrigation Companies in Utah. 73. Snow-Melting Characteristics. *Utah A. E. S. Bull. Nos.* 185, 199, 231 (*Tech.*). 74. Biology and Control of Citrus Insects and Mites. 75. Seasonal Changes in the Chemical Composition of Range Forage and their relation to Nutrition of Animals. 76. Tests of Farm Organization in the Turlock Area. 77. The Composition of Canning Tomatoes. 78. The Vegetable Weevil. 79. Supply, Demand, and Prices of California Peaches. 80. Chemical Control of the Garden Centipede, *Scutigera Immaculata*. 81. The Economic Significance of Tree Size in Western Sierra Lumbering. 82. The Effect of Current Interruption in Electrical Incubation. *California A. E. S. Bull. Nos.* 542, 543, 544, 545, 546, 547, 548, 549, 550. 83. Variation in Citrus Seedlings and their Relation to Root-Stock Selection. 84. Variation in the Yields of Fruit Trees in Relation to the Planning of Future Experiments. 85. Transmission of Carrot, Parsley, and Parsnip Yellows by *Cicadula Divisa*. 86. Injurious Effects of Manganese and Iron Deficiencies on the Growth of Citrus. 87. Plant Buffer Systems in Relation to the Absorption of Bases by Plants. 88. Vaccination of Swine Against Tuberculosis with Calmette-Guerin Culture, BCG. 89. Deterrent Effect of Artificial Light on the Codling Moth. 90. Field Observations on the Beet Leafhopper, *Eutettix tenellus*, in California. 91. The Use of Arsenical Compounds in the Control of Deep-rooted Perennial Weeds. 92. Reproduction without Males in Aseptic Root Cultures of the Root-knot Nematode. Development of the Root-knot Nematode as affected by Temperature. 93. A Comparison of *Aonidiella Aurantii*, and *Aonidiella Citrina*, including a Study of the Internal Anatomy of the latter. 94. The Biology of Bean Thrips. *Hilgardia Vol.* 7, *Nos.* 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12. 95. The Composition of Dairy Products in Six New Jersey Townships. 96. Farm Taxation in New Jersey. 97. Feather Development in Barred Plymouth Rock Chicks. 98. Vireos, Cuckoos, and Shrikes of New Jersey. 99. A Comparison of the Pollution and Natural Purification of the Connecticut and Delaware Rivers and the Brandywine Creek. 100. Factors Affecting the Use of Nitrate and Ammonium Nitrogen by Apple Trees. 101. Analyses of Materials sold as Insecticides and Fungicides during 1932. 102. Analyses of Commercial Fertilizers. Fertilizer Supplies, and Home Mixtures for 1932. 103. Results of Seed and Legume Inoculant Inspection for 1932. 104. Analyses of Commercial Fertilizers and Ground Bone; Analyses of Agricultural Lime. 1932. 105. Proceedings of the New Jersey Land use Conference. 106. Attracting Winter Birds to the Garden and Home Grounds. *New Jersey A. E. S. Bull. Nos.* 506, 532, 543, 544, 545, 547, 548, 549, 550, 551, 552, 553. 107. Mastitis. 108. County and Township Taxes in New Jersey. 109. Spraying Recommendation for Apples. 110. Honey—Its Use in Cookery. 111. Raise your own Food on the Farm. 112. Farm Profits and Factors Influencing Farm Profits on 55 Dairy Farms in Warren County. 113. Factors that Affect the Fat Content of Milk. 114. The Possibilities of Part-time Farming. 115. Minor Enterprises for Part-Time Farmers. 116. Canning Vegetables and Fruits for Winter. *New Jersey A. E. S. Extn. Bull. Nos.* 100, 101, 102, 103, 104, 105, 106, 107, 108, 109.

E. Circulars, Leaflets &c.

117. The Making of Rain-watered Compost Manure from Farm Wastes. *Indore Inst. of Plant Ind. Leaflet No.* 2. 118. Woolly Aphis. *Eng. Mini. Agri.*

& Fish. Adv. Leaf. No. 187. 119. Australian Export Apple Cases. *Australia Conn. Sci. & Ind. Res. Pamph. No. 45.* 120. Frog eye Disease of Tobacco. *Australia Tobacco Investigation Pamphlet No. 2.* 121. United States Grades, Color Standards, and Packing Requirements for Honey. 122. Orchard Insects of the Pacific Northwest and their Control. 123. Effects of Time of Planting and of Fertilizer Mixtures on the Curly-Top Resistant Sugar-Beet Variety U. S. No. 1 in Idaho. 124. The Katahdin and Chippewa Potatoes. 125. The Commercial Storage of Fruits, Vegetables, and Florists' Stocks. 126. Portable Refrigeration Chambers for Studying Cold Resistance of Plants in the Field. 127. Fight Grasshoppers by Plowing Stubble. *U. S. A. Agri. Dept. Cir. Nos. 24, 270, 273, 276, 278, 285, 302.* 128. Improving Cattle in Areas freed of Ticks. 129. Cross Blocking Sugar Beets by Machine. 130. Huron Tomothy. 131. Injury to Buildings by Termites. *U. S. A. Agri. Dept. Leaflet Nos. 51, 97, 99, 101.* 132. Abstracts of Bulletins 442-473, Circulars 61-66 and other Publications during 1932. *Texas A. E. S. Cir. No. 68.* 133. Disinfectants and Disinfection. 134. Care of Milking Machines. *Wisconsin A. E. S. Cir. Nos. 256, 259.* 135. Dry-Farming in Utah. 136. Tomato Culture in Utah. 137. Annual Summary of Publications. *Utah A. E. S. Cir. Nos. 21, 63, 100.* 138. Suggested Method for Treating Chlorosis. 139. Floriculture. 140. Poisoned Bran Bait for Grasshoppers and Cutworms. 141. A New Rabbit Repellent for Protection of Young Orchards. *Utah A. E. S. Mimeograph Sheet. Nos. 23, 43, 55, 73.* 142. State Laws Concerning the Sale of Seeds and Legume Inoculants. 143. Swine Management. 144. Some Poisonous Plants of New Jersey. 145. Facts about 36 Varieties of Peaches. 146. Winter Feeding of Dairy Heifers. 147. Clothes Moths and Carpet Beetles. 148. House Ants. 149. Commercial Fertilizers and Soil Acidity. 150. The Septic Tank. 151. Horse Feeding. 152. Sweet Clover as a Soil Improvement for Crop Orchards. 153. Rapidly Growing, Succulent Branches on Young Apple trees Trend to Form Narrow Crotch Angles with the Trunk. 154. New Jersey Standard for Classifying the set of Fruit Buds upon Peaches. 155. Pyrethrum: Its Value in Exterminating Insects. 156. Rote-none: Its Insecticidal Value. 157. Classification of 135 Varieties and Species of Peaches and Nectarines on Basis of Fruit Bud Set at New Brunswick. 158. Cranberry False Blossom. 159. Movable Hog Houses. 160. Sheep Feeding. 161. Blossom-End Rot of Tomatoes. 162. Rust of Apples. 163. Corn Smut. 164. Establishing a Strawberry Planting. 165. Orange Rust of Blackberry and Raspberry. 166. Cabbage Yellows. 167. Bacterial Wilt of Sweet Corn. 168. Increasing the Vitamin D Content of Milk. 169. Amount of Mulch Material Required by Apple Trees. 170. Elberta and its Selfed and Chance Seedlings Lack Hardiness. 171. The Care of Evergreens. 172. China Asters. *New Jersey A. E. S. Cir. Nos. 248, 259, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289.*

F. Reprints.

173. Contributions to Some Aspect of Gene-Type Analysis of the Garden Bean. 174. Propagation of Equisetum from sterile aerial shoots. 175. The Production of Vestigial and Sterile Sex-Organs through Sex Reversal and Neutral Sexual States. 176. Saving Time and Storage in Breeding Sugar-Beets. 177. Drainage of Land Overlying an Artesian Basin. 178. Some Effects of Wounding Onion Bulbs on Seed Production. 179. Methods of Alleviating Water Shortages on Irrigation Projects. 180. Rainfall and Crop Yields. 181. A Practical System of Soil Diagnosis—Sorghums of all Kinds.

G. Translations. (United States Bureau of Plant Industry.)

182. (1) On Fertilizing Fibre Plants. (2) The Fertilizing Effect of Single Inorganic Materials on Fibre Plants.

DECEMBER 1933

A. Books.

1. The Technique of Artificial Insemination. *Imp. Bur. Animal Genetics Pub.* (1933). 2. Diseases of Plants (A Dissertation). *J. C. Fabricius.* (1926). 3. Observations on the Rust of Grain (1767) tr. by P. P. Pirone. *F. Fontana.* (1932). 4. Taxation of Income in India. *V. K. R. V. Rao.* (1931).

B. Reports.

1. Report on the Operations of the Department of Agriculture, Madras Presidency for the year 1932-33. 2. Report of Subordinate Officers of the Department of Agriculture, Madras for 1932-33. 3. Statistical Tables relating to Banks in India—1931. 4. Index Numbers of Indian Prices—1861-1931. 5. Agricultural Statistics of India—1930-31, Vol. 1. 6. Administration Report of the Department of Agriculture, British Guiana for the year 1932.

C. Bulletins, Memoirs &c.

7. Report on the Manurial Experiments Carried out in Burma by the Agricultural Department from 1912-13 to 1930-31. *Burma Agri. Dept. Bull. No. 29.* 8. The Influence of Climate on Yield Variations. *Imp. Bur. of Soil. Sci. Trans.* 9. Beginnings of Co-operative Grain Marketing. *U. S. A. Federal Farm Board Pub* 10. Co-operative Marketing of Farm Products. 11. Dairy Products: Guide for Organizing Dairy Co-operative Marketing Associations. *U. S. A. Federal Farm Board Bull. Nos. 10, 11.* 12. Increasing the Color of Cranberries after removal from the Vines. *N. Jersey A. E. S. Bull. 504.* 13. Porous Hose Irrigation. *Michigan A. E. S. Extn. Bull. 133.* 14. Price Movements of Agricultural Products: Diagrams of Monthly Gold Prices from 1927 to 1933. *E. M. B. Statistics & Intelligence Branch Pub.* 15. On the Temperature of Atmospheric Layers in the neighbourhood of the Soil. (In French). *Spscl. Pub.*

ADDITIONS TO THE LIBRARY, JANUARY 1934

A. Books.

1. The Trees of New South Wales. *Anderson, R. H.* (19 2). 2. Research in Transportation in relation to Agriculture—Scope and Method. *Black, J. D. Ed.* (1933). 3. Plant Breeding in Soviet Union. *Imp. Bur. Plant Genetics Publication No. 13.* (1933). 4. A Class Book of Botany. *Dutta, A. C.* (1933). 5. Critical Notes on C. D. Darlington's "Recent Advances in Cytology". *Belling, J.* (1933). 6. A Bibliographical Monograph on Plant Genetics--1900-1929 (2nd Edn. Revd. and Enlarged). *Matsuura, H.* (1933). 7. Influence of Manures on the Wilt Disease of *Cajanus Indicus* Spreng and the Isolation of Types resistant to the—Disease (Ind. Sci. Monograph No. 7). *McRae, W. & Shaw, F. J. F.* (1933). 8. The Silk Industry of Japan. (Ind. Sci. Monograph No. 8). *Ghosh, C. C.* (1933). 9. Census of India—Madras Report and Tables. (1 Set). (1932). 10. India in 1931-32. (1933). 11. Alphabetical List of Villages of the Madras Presidency (Corrected up to Sept. 1930). (1933). 12. Madras District Gazetteers—Statistical Appendix to Madura, Bellary and Trichinopoly Districts. (1933).

B. Reports.

1. Madras Agricultural Station Reports for 1932-33. 2. Season and Crop Report of the Madras Presidency for the Agricultural year 1932-33. 3. Annual Report of the Imperial Council of Agricultural Research for the year 1932-33. 4. Report on the Administration of the Meteorological Department of the Government of India in 1932-33. 5. Proceedings of the Twentieth Indian Science Congress, Patna, 1933 (Third Circuit).

C. Bulletins, Memoirs &c.

6. A Collection of Telugu Sayings and Proverbs Bearing on Agriculture. *Madras Agri. Dept. Bull. No. 31.* 7. Testing and Selection of Commercial Wood Preservatives. *Indian Forest Bulletin No. 8—1933.* 8. Lucerne: Its Ecological Position and Distribution in the World. *Imp. Agri. Bur. Herbage Pub. Ser. Bull. No. 12.* 9. Poultry Keeping on the General Farm. 10. The Culture of Fish in Ponds. *England Min. Agri. & Fish. Bulletin Nos. 8, 12.* 11. Rice in British Guiana—1927-1932. *British Guiana Agri Dept. Rice Bull. No. 1.* 12. Physiological Factors affecting the Fruiting of Cotton with Special reference to Boll Shedding. *Arizona A. E. S. Tec'n. Bull. No. 46.* 13. Grain Sorghums in California. *California A. E. S. Cir. 74.* 14. Weeds of the Pineapple fields of the Hawaiian Islands. *Hawaii Univ. Res. Publ. No. 6.* 15. Vegetable Insects and their Control. *Canada Agri. Dept. Bull. 161—New Ser.*

D. Translations.

16. On the Complex of Factors Governing Sexual Reproduction in Millet by Dogushin, D. A. 17. Sorghum by L. Pynaert.

Departmental Notifications.

Gazette Notifications. M. R. Ry. Rao Bahadur T. S. Venkatraman Avl, Soil Physicist, Madras, on other duty under the Government of India to be Deputy Director of Agriculture in the I. A. S. in an existing vacancy with effect from 24th October 1933, in the chain of arrangements made consequent on the retirement of M. R. Ry. Rao Bahadur C. Tadulinga Mudaliar Avl Mr A. Chinnathambi Pillai, Officiating Assistant Director of Agriculture. Guntur, to officiate as Assistant Director of Agriculture, Coimbatore. Mr. A. Gopalakrishnayya Naidu, Upper Subordinate to officiate as Assistant Director of Agriculture, Guntur. Dr A. Subba Rao, M A., M. Sc., D. Sc., is selected for appointment to the temporary post of Soil Physicist. Mr. C. R. Srinivasa Ayyangar, Superintendent, A. R. S. Maruteru, to be special Officer for the study of Rice trade. Mr. K. Venkatrama Ayyar, Superintendent, A. R. S. Pattambi to be Superintendent, A. R. S. Maruteru. Mr. M. C. Cherian, A. D. A. and Offg. Lecturer in Entomology to be Lecturer in Agriculture, without prejudice to his officiating appointment as Lecturer in Entomology.

D. A's Office Orders. Mr. T. K. Mukundan whose offg. appointment as Upper Subordinate, Agricultural section, (Vice Mr. M. C. Menon on leave) terminates on 13th February 1934, will continue to officiate from 14th February till further orders, Vice Mr. A. Chinnathambi Pillai on other duty. Consequent on the reversion of Mr. K. Govindan Nambiar, Upper Subordinate, Agricultural section, III grade to a provisionally substantive post. Mr. V. Satagopan Ayyangar, Upper Subordinate Agricultural Section, IV grade will be provisionally substantive in the IV grade from 3rd January 1934.

Promotions. The following promotions are ordered in the Upper Subordinate service, Agricultural Section, with effect from 13-4-33: from V grade to IV grade:— (1) Mr. P. Gowrisankara Ayyar, (2) Mr. A. M. Muthiah Nattan.

Transfers. Mr. R. Govindaramayya, A. D., from III to V circle. Mr. V. K. Appaji, Upper Subordinate under training at the A. R. S. Aduturai to IV circle as F. M. Palur. Mr. K. Balaji Rao, A. F. M. Hagari, to be A. D. Hospet. Mr. T. Krishna Reddy, A. D., Hospet to be A. D., Gooty.

Leave. Mr. S. Suryanarayana, A. D., Parvatipuram I. a. p. on M. C. for two months from date of relief. Mr. B. N. Padmanabha Ayyar, F. M. Maruter I. a. p. for three months from 1-2-34. Mr. M. P. Narasimha Rao, A. D., Kaikalur, I. a. p. on M. C. for three months from 8-2-34. Mr. S. P. Fernando, A. F. M. Koilpatty, extension of I. a. p. on M. C. for two months from 8-2-34. Mr. V. Panduranga Rao, Assistant, Millets Section I. a. p. for two months from 13-2-34.

The Ramasastrulu—Munagala Prize, 1934.

1. The prize will be awarded in July 1934.
2. The prize will be in the form of a Medal and will be awarded to the member of the Union who submits the best account of original research or enquiry, carried out by him on any agricultural subject.
3. The subject matter shall not exceed in length twelve foolscap pages, type-written on one side.
4. Intending competitors should notify the President or Vice-President of the Madras Agricultural Students' Union not later than the 15th May the subject of the paper which they propose to submit, and the paper should be sent in so as to reach the President or Vice-President, Madras Agricultural Students' Union not later than the 1st June 1934, with a covering letter showing full name and address of the sender. The author's name should not be shown on the paper, but should be entered under a nom de plume.
5. Four type-written copies of the essay should be sent in.
6. The name of the successful competitor will be announced and the prize awarded at the time of the Conference.
7. Papers submitted will become the property of the Union, and the Union reserves to itself the right of publishing all or any of the papers.
8. All references in the paper to published books, reports or papers by other workers must be acknowledged.
9. Any further particulars may be obtained from the President or Vice-President, Madras Agricultural Students' Union, Lawley Road, P.O., Coimbatore.

ADDITIONS TO THE LIBRARY, FEBRUARY 1934

A. Books.

1. Fream's Elements of Agriculture. (12th Edn.). *Biffen, R. H. Ed.* (1932).
2. Farm Tractors. *Stone, A. A.* (1932).
3. Progress in English Farming VII—The Flexibility of Farming. *Bridges, A.* (1933).
4. Financing the Farm. *Orwin, C. S.* (1933).
5. The Causes and Phases of the World Economic Depression. *League of Nations Pub.* (1931).
6. Year Book of Agricultural Co-operation. *Plunkett Foundation Pub.* (1933).
7. Veterinary Obstetrics and Zootechnics. *Beaman, H. N.* (1932).
8. Elementary Histological Technique for Animal and Plant Tissues. *Holder, J. T.* (1931).
9. How Plants get their Names. *Bailey, L. H.* (1933).
10. Evolution, Heredity and Variation. *Cutler, D. W.* (1932).
11. The Pneumatic System of Plants especially Trees. *Macdougal, D. T. & Working, E. B.* (1933).
12. Root Development of Vegetable Crops. *Weaver, J. C. & Bruner, W. E.* (1927).
13. Bacteriology. *Tanner, F. W.* (1933).
14. The Newer Knowledge of Bacteriology and Immunology. *Jordan, E. O. & Falk, I. S.* (1929).
15. Laboratory Outlines in Bacteriology. *Norton, J. F. & Falk, I. S.* (1926).
16. A Text Book of General Bacteriology. *Jordon, E. O.* (1933).
17. Physiology of Bacteria. *Rahn, O.* (1932).
18. Diptera—Vol. IV—Anophalini. (Fauna of British India Series). *Christophers, S. R.* (1933).

B. Bulletins, Memoirs &c.

1. Fruit Survey Report: A Report on Present Condition and Prospects of Fruit Growing Industry in H. E. H. the Nizam's Dominions. *Hyderabad Agri. Dept. Bull. No. 8, 1933.*
2. The Improvement of Grassland.
3. Rations for Live Stock.
4. Cabbages and Related Green Crops.
5. Onions and Related Crops. *Eng. Mini. Agri. & Fish. Bull. Nos. 3, 48, 53, 69.*
6. Investigations on the Standardization of Citrus Trees by Propagation Methods.
7. Problems of Fruit Tree Nutrition: Possible Lines of Approach. *Imp. Bur. of Fruit Production Tech. Comm. Nos. 3, 4.*
8. Fertilization and Crop: Summary of the Results obtained from Field Experiments with Fertilizers. *Imp. Bur. of Occasional Paper No. 2.*
9. Problems of Animal Nutrition and Animal Husbandry in Northern Nigeria. *Imp. Bur. of Animal Nutrition Tech. Comm. No. 4.*
10. The Influence of Root-Stocks on the Texture and Flavour of Orange Fruits. *S. Australia Agri. Dept. Bull. No. 276.*
11. Effects of Stress Conditions on the Cotton Plant in Arizona. *U. S. Agri. Dept. Tech. Bull. No. 392.*

C. Leaflets, Circulars &c.

12. American Tobacco Types: Uses and Markets. *U. S. A. Agri. Dept. Cir. No. 249.*
13. The Kestrel.
14. The Turnip Gall Weevil.
15. Gooseberry Cluster-Cup Rust.
16. Wireworms. *Eng. Mini. Agri. Fish. Adv. Leaf. Nos. 192, 196, 198, 199.*

D. Translations.

17. Vernalization of Agricultural Plants by T. D. Lyssenko (Two Articles).

Departmental Notifications.

Gazette notifications. Mr. S. Dharmalingam Mudaliar, Assistant, Paddy Section, Coimbatore, to officiate as Superintendent, A. R. S. Maruteru, vice Mr. C. R. Srinivasa Iyengar on other duty. The posting of Mr. K. Venkatarama Iyer to Maruteru is cancelled. Mr. C. Vijayaraghavacharya, Assistant, Millet Section, has been posted as Superintendent of the Dairy Farm scheme at Hagari. Mr. R. Swami Rao, officiating Superintendent, Central Farm, Coimbatore, l. a. p. without M. C. for three months from date of relief.

Postings. The following officiating appointments are ordered in the Madras Agricultural Subordinate servic-class I, Upper subordinates, iii grade in the scale of Rs. 75—7½/2—105 with effect from 28th March 1934. (1) Mr. V. Venkatadri Reddi B. Sc. Ag. to officiate as Upper subordinate, Agricultural Section, till further orders, vice Mr. A. Gopalakrishnayya Naidu Garu on other duty, to report to duty at Guntur. (2) Mr. Muhamad Obaidulla Shah B. Sc. Ag., to officiate as Upper subordinate, Agricultural Section till 14th May 1934, vice Mr. K. P. Sankunni Menon on leave, to report to duty at St. Thomas Mount.

Promotions. Mr. S. R. Srinivasa Iyengar, Upper subordinate Agricultural Section, IV grade is promoted to iii grade, provisionally with effect from 28th March 1934—vice Mr. K. Ramanujacharya seconded.

Transfers. Mr. P. S. Athmarama Iyer, F. M. Central Farm, Coimbatore, is transferred to V circle. Mr. S. Sundaram, permanent Assistant, Cotton Section, and temporary Assistant Chemist, in the Madras pempheres and Physiological scheme is transferred to Koilpatty as temporary Assistant in the Madras Fodder Chulam Scheme. Mr. D. Shanmuga Sundaram Pillai, A. D. Central Depot, Madura, is transferred to A. R. S. Koilpatty. Mr. V. K. Kunhunni Nambiar F. M. Nileshwar ii Station is transferred to Kasargod vice Mr. K. W. Chakrapani Marar on leave. Mr. P. K. Kannan Nambiar A. F. M. Kasargod is transferred to Nileshwar ii Station.

Leave. Mr. R. Sabrahmania Iyer, Agricultural Instructor, Borstal school, Palamcottah, is granted l. a. p. for 3 months from date of relief. Mr. V. Viswanatham, A. F. M. Central Farm, Coimbatore, l. a. p. for three months from 10th April or date of relief. Mr. V. Panduranga Rao, extension of l. a. p. for one month from 13th April. Mr. P. S. Narayanaswami Iyer, Assistant, Entomology Section, extension of l. a. p. on M. C. for 6 weeks. Mr. K. W. Chakrapani Marar F. M. Kasargod, l. a. p. for 2 months from 16th April or date of relief. Mr. K. V. Sheshagiri Rao, A. A. D. Allagada, extension of l. a. p. on M. C. for three months and 5 days from 6th March. Mr. K. S. Krishnamurthi Iyer, A. D. Cuddalore, l. a. p. for 2 months from 5th May. Mr. Bhairy Shiva Rao, A. D. Madanapalle, l. a. p. for 2 months from 1st May. Mr. S. Muthuswami Iyer, F. M. Palur, l. a. p. for three months from 15th April. Mr. M. P. Gowrisankara Iyer, A. D. Papanasam, l. a. p. for two months from 16th April.

The Ramasastrulu—Munagala Prize, 1934.

1. The prize will be awarded in July 1934.
2. The prize will be in the form of a Medal and will be awarded to the member of the Union who submits the best account of original research or enquiry, carried out by him on any agricultural subject.
3. The subject matter shall not exceed in length twelve foolscap pages, type-written on one side.
4. Intending competitors should notify the President or Vice-President of the Madras Agricultural Students' Union not later than the 15th May the subject of the paper which they propose to submit, and the paper should be sent in so as to reach the President or Vice-President, Madras Agricultural Students' Union not later than the 1st June 1934, with a covering letter showing full name and address of the sender. The author's name should not be shown on the paper, but should be entered under a nom de plume.
5. Four type-written copies of the essay should be sent in.
6. The name of the successful competitor will be announced and the prize awarded at the time of the Conference.
7. Papers submitted will become the property of the Union, and the Union reserves to itself the right of publishing all or any of the papers.
8. All references in the paper to published books, reports or papers by other workers must be acknowledged.
9. Any further particulars may be obtained from the President or Vice-President, Madras Agricultural Students' Union, Lawley Road, P.O., Coimbatore.

ADDITIONS TO THE LIBRARY, MARCH 1934.

A. Books.

1. Science and the Weather. *Little, W. B.* (1933).
2. Farm Tractors. *Stone, A. A.* (1932).
3. Gardening with Herbs for Flavour and Fragrance. *Helen, M. Fox.* (1933).
4. Sugar Industry and Labour in the United Provinces. *Agarwala, R. D.* (1933).
5. Plantation White Sugar Manufacture, tr. by J. P. Ogilvie. *Harloff, W. H. Th.* (1913).
6. Farm Accounts in the Punjab 1931—32. (Board of Economic Inquiry, Punjab Publication No. 32). *Kartar Singh, S.* (1933).
7. An Economic Survey of Naggal. (Board of Economic Inquiry, Punjab Publication—Village Surveys 5). *Sher Singh.* (1933).
8. An Economic Survey of the Haripur and Mangarh Taluqs. (Board of Economic Inquiry Punjab, Publication No. 9). *B'rai Mul Raj.* (1933).
9. Research in Transportation in relation to Agriculture. *Black, J. D. Ed.* (1933).
10. An Introduction to Geology:—Vol. I—Physical Geology. Vol. II—Historical Geology. *Scott, W. B.* (1932).
11. Plants Useful to Man. *Robins, W. W. & Ramaley, F.* (1933).
12. Gramineae (Maydeae—Paniceae) Flora of Tropical Africa, Vol. IX—Part 6. *Prain, D.* (1934).
13. Index Kewensis—Supplement No. 8. 1926—1930. (1933).
14. Biology in Education. *Crowther, J. G. Ed.* (1933).
15. Recent Advances in the Study of Plant Viruses *Smith, K. M.* (1933).
17. The Senses of Insects. *Eltringham, H.* (1933).
18. Coleoptera Lamellicornia—Fam. Scarabaeidae—Sub. Fam. Euchirinae—Phaenomerinae. (Genera Insatrum Fac. 195). *Ohaus, F.* (1933).
18. Plant Parasitic Nematodes and the Diseases they Cause. *Goodey, T.* (1933).
19. Enzyme Actions and Properties, tr. by R. P. Walton. *Waldschmidt-Leitz, E.* (1929).
20. pH and Its Practical Application. *LaMotte, F. L. Etc.* (1932).
21. Science in the Changing World. *Mary Adams Ed.* (1933).
22. Electrical Year Book—1934. (1934).
23. Mechanical World Year Book—1934. (1934).
24. Gazetteer of the Ramnad District—Vol. III—Statistical Appendix. (1933).
25. Gazetteer of the Guntur District. Vol. III—Statistical Appendix. (1933).

B. Reports.

1. Reports of the Experiment Stations of the Empire Cotton Growing Corporation for 1932—33.

C. Bulletins & Publications.

2. Beekeeping: A Subsidiary Occupation for the Ryots. *Mysore Agri. Dept. Ent. Ser.—Bull. No. 10.*
3. Grassland Research in Australia—*Imp. Bur. Plant Genetics Herb. Plants Bull. 14.*
4. The Manuring of Veg table Crops. *England Mini. Agri. & Fish Bull. No 71.*
5. A Study of Claypan Soils. *U. S. A. Agri. Dept. Tech. Bulletin No 399.*
6. Factors Affecting Experiment Error in Field Plot Tests with Corn. *Iowa A. E. S. Res. Bull. 163.*
7. Notes on the Natural Color Preservation of Philippine Plant—Materials for Museum or Exhibition Purposes. *Philippine Univ. Nat. and App. Soi. Bull.—Vol. III, No. 1, July 33.*

D. Translations.

8. Investigation of the Paddy Seedling Decay in Formosa by Kaneyoshii Sawada.
9. The Influence of Untimely Cold Weather Development of Grain and the Yield in Rice.
10. Full Summary of Karyo-Systematic Investigations in the Family Gramineae by Avdulow, N. P.

UNIVERSITY OF MADRAS

B. Sc. Ag. Degree Examination Question Papers, 1934.

(New Regulations).

FIRST EXAMINATION

1. AGRICULTURE

(Wednesday, 4th April. 7 A. M. to 10 A. M.)

Only six questions are to be answered.

1. Discuss the theory of the formation of deep laterite soils as found on the West Coast, and detail the weathering agencies responsible.
2. What are the effects of the incorporation of large quantities of organic matter on (a) a sandy soil, and (b) a clay soil?
3. Trace the influence of rainfall and climate on the farming practices of the West Coast as against those in the district of Coimbatore.
4. Enumerate the difficulties which may be met with in producing a good deep tilth on a heavy soil as against a light one.
5. Sketch and name the various parts of a mechanical seed-drill.
6. Sketch the essential parts of (a) a Coimbatore district wooden plough, and (b) a double-handled iron plough with beam.
Name the parts and briefly describe the functions of each.
7. What are the factors which influence climate? Indicate their importance.
8. Discuss the importance of weather forecasts to the farmer, and the methods you would adopt so that forecasts reached the farmer in time to be of value to him.

2. BOTANY

(Wednesday, 4th April. 1 P. M. to 4 P. M.)

Only six questions are to be answered. Questions 4 and 7 are compulsory.

1. Describe, with sketches, the structure and the germination of a seed of Lablab, and compare it with that of Maize.
2. Give a brief account of the various kinds of 'Inflorescence', with examples and drawings.
3. Enumerate, with sketches, the various kinds of tendrils you have examined. Explain of what part of the plant each is a modification, and how each functions.
4. Write a short note on the morphology of cabbage. Explain why the leaves of the 'heart' are of lighter colour, sweeter taste, and more tender texture than the external ones.
5. What are meristematic tissues? Give an account of those found in a dicotyledonous stem.
6. Explain the changes that take place in the nucleus when a cell undergoes division.
7. Mention the important characteristics of the families, Solanaceae and Cucurbitaceae, and describe botanically any well-known plant in either of them.
8. Write short notes on:—Cladode, Syconium, Phellogen, Cystolith, Calypotropen, Velamen, Cyathium, Vivipary, Synergidae, and Thyrsus.

3. CHEMISTRY

(Thursday, 5th April. 7. A. M. to 10 A. M.)

Only six questions are to be answered.

1. Explain, with examples, (a) saturated and unsaturated compounds, (b) open and closed chains, and (c) constitutional and molecular formulae.

2. (a) How would you prepare a specimen of iodoform in the laboratory? What happens when it is boiled with caustic soda?

(b) How would you ascertain the purity of a sample of chloroform?

3. Describe the preparation of acetone from calcium acetate, and show what light this reaction throws on the constitution of acetone. What are the characteristics of the ketones as a class?

4. How is glycerine prepared on a large scale? Give its important properties and its uses in industries.

5. A substance isolated from a plant has the empirical formula CHO_2 and its molecular weight is 90. It is an acid and forms soluble salts with potassium, and sodium, and insoluble salts with calcium and lead. It is completely oxidized by potassium permanganate in acid solution, but is unaffected by dilute nitric acid. What are the conclusions you would draw from these facts as to its nature? What would be the effect of strong sulphuric acid on this substance?

6. What are the various forms of lactic acid known? How is their existence explained? Give the sources from which they can be prepared.

7. Describe any method of preparing methylamine. Show how amines resemble ammonia. Indicate how primary, secondary, and tertiary amines can be distinguished.

8. Explain the meaning of the term 'phenol'. Describe the preparation of 'carbolic' acid from coal tar, and explain why the substance is called an acid.

4. ZOOLOGY

(Thursday, 5th April. 1 P. M. to 4 P. M.)

Only six questions are to be answered.

1. What do you understand by the terms lower and higher animals? What are their chief similarities and differences? Explain, with the help of diagrams, the main differences in structure between a cockroach and a rabbit.

2. Describe how the following animals perform the function of respiration:— a crab, a grasshopper, a spider, a fish, and a bird.

3. Classify the following animals into their respective groups, and state in a line or two, what you know of each:— cuttle-fish, guinea-worm, chameleon, leech flying fox, turtle, tick, oyster, *Peripatus*, and *Opalina*.

4. Explain clearly what you understand by the terms:—abiogenesis, bilateral symmetry, mimicry, natural selection, and metamorphosis.

5. Describe briefly the structure and arrangement of the organs of an insect concerned in the act of feeding, pointing out the main differences found among different kinds of insects.

6. Discuss the important points in the homologies of the body segments in:— a prawn, a spider, a centipede, and a grasshopper.

7. Write a brief account of parasitism found in the animal kingdom, pointing out its effect on the individual and its economic importance.

8. Write short notes on:—termites, jellyfish, tapeworm, millipedes, and larva

(New Regulations)

SECOND EXAMINATION

1. AGRICULTURE—PLANT HUSBANDRY I

(Wednesday, 4th April. 7 A. M. To 10 A. M.)

Only five questions are to be answered. Question 3 is compulsory.

1. What is compost? How does it differ from poudrette? Explain how both are made and state to what crops, and in what quantities they can be applied.

2. What are rotations and mixtures? In what ways do they differ? What are their relative advantages?

3. From your experience of plot cultivation at the College farm, which type of land would you prefer to have—dry, garden, or wet? Give reasons for your choice. How many acres of the type of land you choose would you need to keep you in reasonable comfort? What would you expect to get out of it per annum?

4. What are the various improvements that you can suggest in the cultivation of any two of the following:—coco-nut, potato cotton and ragi?

What advantages will these improvements give to the ryot adopting them?

5. What is the difference between hay and silage? Describe how either is made, and estimate the cost of making 10 tons of the same.

6. What are the ingredients in the following manures; to what crops can they be applied, and in what doses; what is the price of each per ton:—bonechar super, niciphos, muriate of potash, superphosphate, castor cake, nitrolim and guano?

7. The sources of irrigation for sugar-cane are a well, a tank, and a channel. What are the irrigation charges in each case till the crop matures?

1. AGRICULTURE. PLANT HUSBANDRY II

(Wednesday, 4th April. 1 P. M. To 4 P. M.)

Only five questions are to be answered. Question 2 is compulsory.

1. How are alkaline soils formed? Discuss briefly the methods of reclamation.

2. Classify the three most important oil-seeds in the Madras Presidency. Detail methods and cost of cultivation, and profits, of the most important.

3. Five tons of unrotted paddy straw were ploughed under in a field of garden land on the Central Farm, Coimbatore. How would the application affect the growth of (a) cholam, and (b) sunnhemp, sown immediately after the straw was ploughed under?

4. Discuss the advantages and disadvantages of bare fallows in crop production.

5. Discuss the economic importance of any one of the following hill crops:—tea, coffee, and potato.

6. Make up a mixture of concentrated manures, showing the amounts of each you would use, to give the following amounts of plant food per acre:—60 lb. of nitrogen (half as nitric nitrogen and half as organic nitrogen); 54 lb. of soluble phosphoric acid; and 50 lb. of potash.

7. A well has just been dug in the corner of a six-acre block of sandy soil. Illustrate and discuss the lay-out of permanent and temporary channels which you would adopt, so that losses by seepage will not be too high and the cost of preparing the channels not too heavy.

(Old and New Regulations.)

SECOND EXAMINATION (New) & PART I (Old)

(Common to Part I under Old Regulations and Second Examination under New Regulations)

2. AGRICULTURAL ENGINEERING

(Thursday, 5th April. 7 A. M. to 10 A. M.)

Only six questions are to be answered. Question 2 is compulsory.

1. (a) Describe briefly an indigenous water-lift with which you are familiar, and determine its efficiency.

(b) Work out the cost of raising 1,000 gallons of water through a height of one foot, by means of the lift described by you. (Assume any values you require for the purpose, but state your assumptions clearly.)

2. It is proposed to instal a 16 h. p. horizontal oil engine on a well in the Coimbatore district for pumping water for irrigation. Design a suitable *engine house* and furnish necessary dimensioned sketches for the same, so that any ryot could build it to your design.

3. Prepare an estimate of cost for the *roof of the engine house* in Question 2.

4. What is the function of a governor in an engine? Explain clearly, with sketches, how it works. Can an engine be worked without a governor? Give reasons for your answer

5. What are the important qualities that a lubricant should possess? What determines mainly the choice of a lubricant for a given purpose? Could groundnut oil be used for lubricating the internal parts of an oil engine? Explain your answer.

6. A plot of land is in the shape of a quadrilateral *ABCD*. Length of *AB* is 400 yds., *BC* is 650 yds., *CD* is 1,405 yds., *DA* is 920 yds., and *AC* is 2,580 ft. Find its area in acres.

7. Describe briefly the Dumpy Level or any other level with which you are familiar, and explain clearly how you would test and adjust it for field work.

8. What is meant by the term 'water table'? What are the causes which affect the water table in any locality?

3. AGRICULTURAL ZOOLOGY

(Thursday, 5th April. 1 P. M. to 4 P. M.)

Only six questions are to be answered. Question 1 is compulsory

1. What are social insects? Give a brief summary of the life-history and habits of any social insect you know, putting out its economic importance, if any.

2. What do you understand by the term 'animal tropisms'? Describe briefly any such tropisms you know, prevalent among insects, and discuss their economic importance, if any.

3. Explain clearly, with the help of examples, the meaning of the following terms when used in connexion with insects:—protective adaptation, ecdysis, fumigation, myiasis, locust, entomophaga, hypermetamorphosis, and hibernation.

4. Classify the following forms into their respective groups or families, and explain the economic importance, if any, of each:—sawfly, mealworm, *Chrysopa* cricket, bollworm, *suval-puchi*, tick, ladybird, eriworm, and Thrips

5. Describe the more important features in the life-histories of the following animals:—the ox bot blister beetle, malaria germ, potter wasp, and the liver fluke.

6. Write what you know of the biological methods of pest control, with special reference to South India.
7. Write brief notes on :—strepsiptera, trap crops, galls, cut-worms, pebrine, soorai, white grubs, and eel-worms.
8. Give a very brief account of the important insects affecting the castor plant in South India under the following heads :—
 - (a) name and family of insect ;
 - (b) nature and extent of damage done ;
 - (c) alternate food plant ;
 - (d) natural enemies ; and
 - (e) control measures you would suggest.

4 ANIMAL HYGIENE

(Friday, 6th April. 1 P. M. to 4 P. M.)

Only six questions are to be answered. Questions 4 and 8 are compulsory.

1. Enumerate the ferments contained in the pancreatic juice of the ox, and discuss the role of each in digestion.
2. Give the actions, uses, and doses of the following drugs in cattle:—
(a) *nux vomica*, (b) turpentine, and (c) eucalyptus oil.
3. Describe the symptoms and treatment of any three of the following:—
(a) spirochaetosis in the fowl, (b) ringworm in the calf, (c) dislocation of the patella in the ox, and (d) parasitic bronchitis in the sheep.
4. Mention the diseases that are communicable to man from farm animals, and, in respect of any two of them, explain in detail how you would arrive at a definite diagnosis.
5. Describe either rinderpest, or black-quarter, as regards aetiology, animals susceptible, mode of infection, methods of diagnosis, and immunology.
6. Describe the rumen of the ox, and define its position in the abdominal cavity.
7. What are the objects of castration? Describe the method of castration you would prefer for cattle in the field, indicating your reasons for the choice.
8. Enumerate the common ailments of the cow, incidental to parturition, and mention briefly under each how you would deal with it

(Old Regulations)

PART I

1. AGRICULTURAL BOTANY

(Wednesday, 4th April. 7. A. M. to 10 A. M.)

Only six questions are to be answered. Question 4 is compulsory.

1. Draw rough sketches and describe (a) a mango fruit, (b) an inflorescence of a coco-nut palm, (c) a potato-tuber, and (d) a radish.
2. Contrast the structure of a green leaf and a mature seed.
3. Describe how the mineral nutrients and water are absorbed by plants.
4. Explain the process of respiration in plants, and describe an experiment to illustrate the same.
5. Write short notes on :— (a) enzymes, (b) cytoplasm, (c) karyokinesis, and (d) mesophytes.
6. State the ecological factors which govern the distribution of plants, and give examples to show how these factors operate.

7. (a) What are the general principles on which the systematic classification of plants is based?

(b) To what families do the following plants belong :— coffee, aloe, potato, and betel?

8. Give the distinguishing characteristics of any two of the following :— Compositae, Leguminosae, Graminaceae, and Euphorbiaceae.

2. AGRICULTURAL CHEMISTRY

(Wednesday, 4th April. 1 P. M. to 4 P. M.)

Only six questions are to be answered. Questions 1 and 8 are compulsory.

1. How would you establish the constitutional formula of acetic acid?

2. Describe the general methods available for preparing aldehydes, and give the important reactions characteristic of the aldehyde group.

3. Write short notes on any four of the following :— (a) isomerism, (b) optical activity, (c) fractional distillation, (d) diazo reaction, (e) substitution product, and (f) unsaturation.

4. What are the products of wood distillation? How would you prepare commercial and pure methyl alcohol from the distillate?

5. What do you understand by the following terms with reference to soils :— saturation capacity, hygroscopic moisture, retention of bases and pore space?

6. What are the differences in the physical, chemical, and biological properties of sandy and clay soils?

7. Give in outline any two methods of the mechanical analysis of a soil, mentioning the principles on which they are based. Of what use is the mechanical analysis of a soil?

8. State briefly the biological processes connected with the breakdown of carbohydrate material in the soil under aerobic and anaerobic conditions.

3. AGRICULTURE, I

(Friday, 6th April. 7 A. M. to 10 A. M.)

Only five questions are to be answered. Question 3 is compulsory.

1. Classify the soils of the Madras Presidency. Discuss their origin. What are the peculiarities of each type?

2. Name the several instruments used for meteorological observations at the Agricultural College and Research Institute at Coimbatore. How are these observations likely to be of use to the educated farmer?

3. In a farm located in a black cotton soil district, a rainfall of 4 in. is received during the third week in August. According to the cropping scheme, two fields, each 5 acres in extent, are to be sown separately with sorghum and cotton. Describe the steps you will take to get the fields ready for sowing, indicating the various operations to be done till sowing is completed. Write out a cultivation sheet for each field, indicating the dates on which each operation is performed, with costs.

4. Give examples of any cases where the ryots of this Presidency follow a systematic rotation of crops. In regard to your own native district, describe, by means of specific instances, how rotations can be profitably adopted by the ryot, and state the advantages he will derive by doing so.

5. Comment on the truth of the statement that canal irrigation is not an unmixed blessing in this country, giving reasons for your opinion.

6. (a) State the effect of the following operations on black soils :—
 - (i) ploughing deep too frequently ;
 - (ii) ploughing soon after a heavy rain ;
 - (iii) applying tank silt at 200 cartloads per acre ;
 - (iv) sowing after half an inch of rain.
 (b) What will be the effects of the above operations on sandy loams ?
7. Describe the methods of reclaiming land under the following conditions:—
 - (a) out of cultivation for ten years ;
 - (b) impregnated with salts injurious to vegetation ;
 - (c) foul with deep-rooted weeds.

PART II

1 AGRICULTURAL BOTANY, I

(Saturday, 7th April. 7 A. M. to 10 A. M.)

Only five questions are to be answered. Questions 3 and 7 are compulsory.

1. Write what you know about the origin of cultivated plants.
2. Give an account of the typical adaptations of plants to conditions of drought, both physical and physiological.
3. Describe in botanical terms *Arachis hypogaea*, and trace the development of its fruit. Illustrate your answer with diagrams.
4. What are weeds? Mention three indigenous and three introduced, troublesome weeds, explaining how they spread, and suggesting means of control.
5. Classify the oil-yielding plants grown in South India, noting their families and explaining from what part of the plant the oil is obtained, and how.
6. Now that prickly pear is being destroyed by the cochineal insect, what suitable substitutes of live fence would you suggest to a farmer to protect his crops, and how would you propagate them?
7. Hill roses do not ordinarily thrive in the plains. With your knowledge in gardening, what methods would you adopt to get them established? Describe the process fully in each case.

1. AGRICULTURAL BOTANY, II.

(Saturday, 7th April. 1 P. M. to 4 P. M.)

Only five questions are to be answered. Question 4 is compulsory.

1. Write short notes on:—(a) segregation and recombination of factors, (b) vegetative and sexual reproduction, and (c) linkage.
2. Explain the various methods of plant breeding which you would adopt, with a view to improve the yields of the following crops in the shortest possible time:—(a) limes, (b) potatoes, (c) paddy, and (d) cumbu.
3. Name the organisms causing the following diseases, and describe the methods of combating them:—(a) paddy-blast, (b) cholera-smut, (c) groundnut-wilt, and (d) coconut bud-rot.
4. An average cultivator wants to improve the yield of groundnuts. Would you advise him to practise 'mass-selection' or 'pure-line-selection'? Give reasons in support of your answer.
5. Explain, with suitable examples, what is meant by the alternation of generations in plants.
6. Explain, with examples, how the study of the life-history of a fungus helps you in the control of plant diseases.
7. Explain briefly the modes of reproduction in—(a) green algae, (b) Characeae, and (c) ferns.

2. AGRICULTURAL CHEMISTRY, I.

(Monday, 9th April. 7 A. M. to 10 A. M.)

Only six questions are to be answered. Question 1 is compulsory.

1. Mention the important contributions made to Agricultural Chemistry by (a) Liebig, (b) Pasteur, and (c) Lawes and Gilbert.
2. What do you understand by the terms 'flocculation' and 'deflocculation' as applied to soils? How are these brought about by farming operations?
3. Explain fully the following terms:—(a) tilth, (b) soil acidity, (c) minimum cropping value, and (d) limiting factors, as applied to manuring.
4. What is meant by alkaline soils? How are they formed, and how would you ameliorate them?
5. How would you estimate the total nitrogen in a sample of castor cake? Explain fully the chemical changes in the several steps involved in the estimation.
6. Describe the sources and properties of sulphate of ammonia and nitrate of soda, and indicate what considerations, apart from their market values, would influence your decision as to which of them should be used on (a) a clay loam, (b) a garden soil, (c) a swampy, paddy soil, and (d) a sandy loam.
7. Name the important indigenous concentrated organic manures available in South India, with the percentages of their predominant manurial constituents.
8. Explain fully the various changes that take place during the storage of farmyard manure.

2. AGRICULTURAL CHEMISTRY, II

(Monday, 9th April. 1 P. M. to 4 P. M.)

Only six questions are to be answered. Question 8 is compulsory.

1. What are essential oils? How are they classified? Describe any method of extracting essential oils. Give the properties of any two of them.
2. What is diastase? Where is it found? How is it extracted? What is its function? Give the chemical changes brought about by the action of diastase on starch.
3. (a) What is, in general, the percentage of albuminoids in leguminous seeds?
(b) How do oil-seeds differ from other seeds? Name one oil-seed, with the percentage of oil contained therein, from each of the following classes:— seeds yielding drying oil, non-drying oil, and semi-drying oil. Give the uses to which these oils are put.
4. Give a short account of starch in food, from the time it is taken up in the mouth of an animal till it is finally disposed of. Mention the chemical changes it undergoes and the agencies which bring about these changes.
5. (a) What are the special requirements of the food of young growing animals as compared to grown-up animals?
(b) Calculate the *total energy* present in the following foodstuffs;—

	Wheat bran (per cent.)	Maize silage (per cent.)
Ether Extract	4.00	2.90
Albuminoids	14.43	8.56
Soluble Carbohydrates ...	57.02	43.14
Fibre	11.85	30.40

6. (a) What is the 'utility of feeding standards'? Give the principles underlying any one of the feeding standards you have studied.

(b) How is the digestibility co-efficient of a foodstuff determined ?

(c) How would you find out the *net available energy* of any foodstuff ?

What information would you require in this calculation ?

7. (a) Give the effects of (i) lactic ferment, and (ii) butyric ferment on milk.

(b) What is sterilization of milk ? How is it done ? What is its effect on milk and the quality of milk ?

8. Discuss the use of preservatives in butter. Mention the preservatives commonly known to be used in butter. What is the effect of added common salt on butter ?

3. AGRICULTURE I

(Tuesday, 10th April. 7 A. M. to 10 A. M.)

Only six questions are to be answered. Questions 3 and 7 are compulsory.

1. Discuss the expenditure in bringing an acre of coco-nuts into bearing and the probable profits when they are in full bearing.

2. For what purposes is seed selection practised, and what are the two main methods ?

3. (a) What are the important points to be looked for in a dairy cow ?

(b) How would you feed a cross-bred dairy cow for 4 gallons of daily milk production ?

4. Discuss the relative advantages of hay and silage in (a) preparation, and (b) feeding to live stock.

5. What kinds and amounts of manures would you advise a farmer to apply as a substitute in the place of 10,000 lb. of farmyard manure ?

6. What difficulties does the small farmer face in maintaining high quality cows and in breeding his own bullocks from them ? Suggest methods of overcoming them.

7. (a) what is meant by 'duty of water' ? What is the duty when 2 inches depth of irrigation is required every seven days ?

(b) Give in detail calculations to show the area which 6 mchotes will irrigate to a depth of $1\frac{1}{2}$ inches in eight hours under Central Farm conditions.

8. Discuss the economics of seed production of the following green manure crops in any district you are familiar with, and explain in detail under what conditions you would grow them :—(a) daincha, (b) sunnhemp, and (c) kolinji.

3. AGRICULTURE, II.

(Tuesday, 10th April. 1 P. M. to 4 P. M.)

Only five questions are to be answered. Question 3 is compulsory

1. What is meant by 'experimental error' ? What steps would you take to minimize it ? What part does randomization play in modern experimentation, and what are its merits and demerits ?

2. The crop specialist gives you three strains of cotton which, he considers, are an improvement on the ryot's crop. What stages should these strains pass through before you can expect any one of them to replace the local variety ?

3. In the neighbourhood of the College Farm there is a holding of 250 acres consisting of 100 acres of dry, 50 acres of wet, and 50 acres of garden land. In which cropping normal to the locality is followed. What is the equipment in permanent labour and in live and dead stock needed for running the farm, and what is the capital required for wages and maintenance for one year ? Explain how you would distribute human and animal labour, and the methods you would employ to maintain the farm most efficiently.

4. Calculate the cost of cultivation of, and the gross and net returns obtainable from, any two of the following crops:—sugar-cane in Vizagapatam, chillies in Guntur, paddy in Malabar, tobacco in Madura, gogu in Godavary, and sunnemp in Tinnevely.

5. Discuss the facilities which exist in South India for marketing any one of the following:—cotton, rice, fruits, and eggs. What, in your opinion, should be the lines along which improvements could be effected to enable the producer to derive the greatest benefit?

6. What are crop cutting experiments? What use can you make of them? Describe how they should be conducted, taking paddy as an example.

7. Give some examples of certain products which the agriculturist now calls waste and which can be utilized for his benefit. Describe how they can be made useful and at what cost.

4. AGRICULTURE—ESSAY

(Wednesday, 11th April. 7 A. M. to 10 A. M.)

Write an essay on:—

Either, The forces that contribute to the increase and decrease of national food-supply.

Or, What developments in agricultural science and agricultural practice, including production and marketing, do you consider of greatest importance to agriculture in India at present?



The Ramasastrulu—Munagala Prize, 1934.

1. The prize will be awarded in July 1934.

2. The prize will be in the form of a Medal and will be awarded to the member of the Union who submits the best account of original research or enquiry, carried out by him on any agricultural subject.

3. The subject matter shall not exceed in length twelve foolscap pages, type-written on one side.

4. Intending competitors should notify the President or Vice-President of the Madras Agricultural Students' Union not later than the 15th May the subject of the paper which they propose to submit, and the paper should be sent in so as to reach the President or Vice-President, Madras Agricultural Students' Union not later than the 1st June 1934, with a covering letter showing full name and address of the sender. The author's name should not be shown on the paper, but should be entered under a nom de plume.

5. Four type-written copies of the essay should be sent in.

6. The name of the successful competitor will be announced and the prize awarded at the time of the Conference.

7. Papers submitted will become the property of the Union, and the Union reserves to itself the right of publishing all or any of the papers.

8. All references in the paper to published books, reports or papers by other workers must be acknowledged.

9. Any further particulars may be obtained from the President or Vice-President, Madras Agricultural Students' Union, Lawley Road, P. O., Coimbatore.

ADDITION TO THE LIBRARY APRIL 1934

A. Books.

1. Meteorology. *Piston, D. S.* (193). 2. The Art of Flower Arrangement in Japan. *Sadler, A. Z.* (1933). 3. Research in Marketing Farm Products—Scope and Method. *Black, J. D., Edn.* (1932). 4. A survey of the Marketing of Fruit in Poona. *Gadgil, D. R. and Gadgil, V. R.* (1933) 5. Principles and Practice of Farm casting with Farm studies. *Patil, P. C.* (1933). 6. The Course and Phases of the World Economic Depression—revised edition. *League of Nations Publication.* (1931). 7. Socialists Planned Economy in the U. S. S. R. (1932). 8. Methods of Statistical Analysis in the Social Sciences. *Davis, G. R. and Crowder, W. F.* (1933). 9. Chemical investigations of the Tobacco Plant. *Vickery, H. B., etc.* (1933). 10. Our changing world view. *Smuts, J. C., etc.* (1932). 11. The Living Universe. *Young Husband, F.* (1933). 12. Classified Catalogue Code. *Ranganathan, S. R.* (1934).

B. Reports.

1. Annual Report of the Indian Lac Research Institute for 1932-33. 2. Mysore Agricultural Calendar for 1934. 2(a). Report for the British Association for the advancement of science—1933. 3. Annual Report of the Agricultural Department, British Guiana for 1932. 4. Annual Report of the Agricultural Department, Palestine for the years 1931 and 1932

C. Bulletins.

1. Preliminary Studies of important crops in the Bombay Deccan in the Post-war Period. *Bombay Agri. Deptt. Bull.* 168. 2. Fruit Productions: Soft Fruits, *Min. of Agri. & Ministry Bull.* 4. 3. Soil, Vegetation and Climate. *Imp. Bur. of Soil Science Tech. Commn.* No. 29. 4. Soils and Fertilisers. *Reprint: Reports of the Progress of Applied Chemistry, Vol. XVIII, 1933.* 5. Practical Kiln-Drying. *Deptt. of Scientific and Industrial Research Forest Products Research Special Report No. 3.* 6. Citru Fruit Prospects in Nyasaland. *Deptt. Agri. Nyasaland Protectorate Bull.* No. 8 (new Ser.) 7. Economic investigation into Livestock Farming in the Northern Transvaal, 1927-30; Part II, the Lowveld Area. *Deptt. Agri. Union of S. Africa; Econ. Series No. 18.* 8. Effect on subsequent yields of storing cut seed potatoes at different temperatures and humidities. *U. S. Deptt. Agri. Tech. Bull.* No. 394. 9. The world situation in oils and oilseeds. *Foreign Agri. Ser. Washington. Reprint.*

D. Leaflets.

Nitrogenous Fertilizers. *Min. of Agri. and Fisheries Adv. Leaf. No. 111.*

ADDITIONS TO THE LIBRARY, MAY 1934

A. Books.

1. World's Grain Exhibition and Conference—Ottawa—Canada—Vol. I, Proceedings. (1933)
2. The Vegetable Products of Ceylon. *Lewis, F.* (1934).
3. The Toxicology of Plants in South Africa. *Steyn, D. G.* (1934).
4. The Common Trees of New Jersey. *Ulrich, J. S.* (1929).
5. Farm Accountancy—Statistics for 1929—1930. *Inter. Inst. of Agri. Pub.* (1933).
6. Chinese Farm Economy. *Buck, J. L.* (1933).
7. Farm Tenantry in the United States. *Bizzoli, W. B.* (1921).
8. Intermediate Book Keeping—5th Edition. *Subramania Ayyar.* (1930).

B. Reports.

1. Administration Report of the Madras Co-operative Department for 1932—1933.
2. Report on Public Instruction in the Madras Presidency for 1932—1933
3. Official Year Book of the Commonwealth of Australia—No. 26, 1933.

C. Memoirs, Bulletins and Special Publications.

4. Famine in the Madras Presidency. *Mad. Govt. Pub.* (1934).
5. The Indian South West Monsoon and the Structure of depressions associated with it.
6. On the Physical Characteristics of Fronts during the Indian South West Monsoon. *Mem. Int. Met. Dep. V, 26/Parts 2, 3.*
7. Tree Planting Book. *Amer. Tree Assn. Pub.* (1922).
8. The Forestry Primer. *Amer. Tree Assn. Pub.* (1933).
9. Bibliography of Helminthology, for the year 1932. *Imp. Bur. of Agri. Parasit. Pub.* (1934).
10. Denudation and Soil Erosion in Nyasaland by A. J. W. Horney. *Nyasa Agri. Dep. Bull. 11.* (1934).
11. The Selection of Citrus Trees, its importance and Problems. *Palestine Agri. For. & Fish. Ser. IV, No. 25.* (1934).
12. Farm and Home drying of Fruits and Vegetables.
13. Diseases of Cabbage and Related Plants.
14. Important Sugar-beet by-Products and their utilization.
15. Improving the Farm Environment for wild life.
16. Judging Sheep.
17. Diseases and parasites of Poultry.
18. Adobe or sun-dried brick for Farm Buildings. *Farmers' Bull. Nos. 984, 1439, 1718, 1919, 1199, 1652, 1723, U. S. Agri. Dep.* (1934).
19. Artificial Drying of Rice on the Farm. 20. Windrowing qualities of Co 231 and other varieties of sugarcane under Louisiana Conditions.
21. Farm Real Estate Situation, (1932—1933). *Cir. Nos. 292, 304, 309, U. S. Agri. Dep. Wash.* (1933).
22. Pollination and Blooming habits of the Persian walnut in California. *Tech. Bull. 387, U. S. Dep. Agri. Wash.* (1934).
23. The Flow of Water in Flumes. *Tech. Bull. 393, U. S. Dep. Agri. Wash.* (1933).
24. Effect on Subsequent yields of storing cut seed potatoes at different temperatures and humidities
25. Hydrion concentration changes in relation to growth and ripening in fruits. *Tech. Bull. Nos. 394, 403, U. S. Dep. Agri. Wash.* (1934).
26. The Comparative Value of different colonies of bees for fruit pollination. *Cornell Agri. Exp. Stn. Memoir No. 117.* (1931).
27. Social and Economic Areas of Broome County, New York. 1928. *Cornell Agri. Exp. Stn. Bull. No. 559.* (1933).
28. Economic Studies of Dairy Farming in New York XI.
29. Studies in Plant Propagation.
30. Correcting the unproductiveness of Acid and Alkaline Muck Soils for the Growing of Vegetable Crops. *New York Agri. Exp. Stn. Bull. Nos. 562, 571, 572.* (1933)
31. The Relative Vigor and Productivity of Potato Plants from Basal and apical sets. *New York State Agri. Exp. Stn. Bull. No. 633* (1933).
32. Lysimeter Investigations. *New York State Agri. Exp. Stn. Tech. Bull. 212.* (1933).
33. The effects of Mulched and Turned Rye in the Green and Mature Stages on the Liberation of Plant Nutrients from a Silt Loam Soil. *Vir. Polytech. Instt. Tech. Bull. No. 53.* (1934).
34. Economic Aspects of the Grade and Staple length of Cotton produced in Oklahoma. *Oklahoma Agri. Exp. Stn. Bull. No. 212.* (1933).
35. Moisture and Fertility Relations

of Subsoil Variations in Heavy Silt Loam Soil at Goodwell, Oklahoma. 36. The Economy of Soil Nitrogen under Semiarid Conditions. 37. Raw Organic Matter Accumulations under various systems of culture. 38. Effect of Saline and Alkaline waters on domestic animals. *Oklahoma Agri. Mech. & Exp. Coll. Bull. Nos. 214, 215, 216, 217.* (1933). 39. Irrigation Investigations in Young Grapefruit orchards on the Yuma Mesha. *Arizona Agri. Exp. Stn. Tech. Bull. No. 37.* (1931). 40. On the Genetic Nature of Induced mutations in Plants. *Missouri Agri. Exp. Stn. Res. Bull. No. 204.* (1933). 41. Varieties of Rice for Texas, *Texas. Agri. Exp. Stn. Bull. No. 485.* 42. Farm Re-organisation and Management. *Dakota Agri. Exp. Stn. Bull. No. 273.* (1933).

D. Reprints.

43. The Effect of Nitrate Supply on Transpiration Ratio in Plants by L. A. T. Ballard: (Reprint from *Aust. Jl. of Exp. Bio. & Med. Sc.* Vol. XI, '33).
 44. The Bacteriological Oxidation of Sulphur in an Australian Soil, by Phyllis M. Rountree (Reprint, Do.).

ADDITIONS TO THE LIBRARY, JUNE 1934.

A. Books.

1. Introduction to Meteorology (Reprint). *A. K. Das & B. N. Srivasthava*. (1933). 2. Weed Destruction by Sulphuric Acid Spraying. *R. K. MacDowall*. (1933). 3. Diptera, Volume V, Fam Culcidae: Tribes—Megartinini and culicini. *P. J. Barraud*. (1934). 4. The Inequality of Man. *J. B. S. Haldane*. (1932). 5. Indian Year Book (1934-'35). (1934). 6. Agricultural Statistics of India, (1930-'31); Volume II—Indian States. (1934). 7. Gazetteer of Tinnevely District, Statistical Appendix—Volume II. (1934). 8. Gazetteer of the West Godavari District; Statistical Appendix—Vol. II. (1934). 9. Anglo-Malayalam Dictionary—Revised and Enlarged. *Zacharias, O. F. E.* (1933). 10. Malayalam—English Dictionary—2nd Edition—Revised and Enlarged. *Zacharias, T.* (1921).

B. Reports.

1. Annual Report of the Agricultural Department, Ontario for 1932. 2. Annual Report of the Agricultural Experiment Station, Massachusetts for 1933.

C. Bulletins, Memoirs, etc.

3. The Sanitary Disposal and Agricultural Utilisation of Habitation Wastes by The Indore Process. *Indore. Instt. Plant. Ind Bull. 1.* (1934). 4. The Coffee Stem Borer *Mys. Agri. Dep. Ent. Bull. No. 11.* (1934). 5. Agricultural Statistics of India, (1930-'31); Volume II, 1934. *Ind. Commer. Intell. & Statistics Dep.* (1934). 6. Note on the Outturn and Profit from Sugarcane Crops grown in the Deccan Canal Tracts. 7. Report on Land Drainage Experiments at Baramati. 8. Note on Research Work done by the Special Irrigation Division since its formation in June, 1916. 9. A Critical Study of Runoff and Floods of Catchments of the Bombay Presidency with a short note on Losses from Lakes by Evaporation. 10. Standard Type Buildings for the Forest Department, Bombay Presidency—Abstracts, Estimates, Block Plans, Specifications, &c. 11. Note on Cement Concrete roads in the United Provinces and on the Financial Aspects of Cement Concrete and asphalt Surfaces. 12. Note on Boring Operations in the Bombay Presidency. 13. Concise Report on the Punjab Irrigation Works. 14. Note on Silt Exclusion from Canals; Part I—Note of May 1931 on Silt Exclusion from Canals—"Silt Pond" versus "Open Flow". 15. Note on Silt Exclusion from Canals—Part II—First Report on Experiments with the full width model of the Sukkur Barrage at Khadakrasla with Appendices on the Design of dynamically similar Models in accordance with Lacey's Formulas. 16. Report on Reclamation and Drainage of Alkali Lands in Hungary. 17. Development of Irrigation in the Deccan Canal Areas. *Bombay P. W. D. Tech. Paper Nos. 21, 24, 28, 30, 37, 40, 41, 43, 45, 46, 48, 49.* 18. Ducks and Geese. 19. Rex-Furred Rabbits. 20. Poisonous Plants on the Farm. *Eng. Min. Agri. & Fish. Bull. Nos. 70, 73, 75.* (1934). 21. Passion Fruit Culture on the North Coast. *N. S. Wales Agri. Dep. Farm. Bull. No. 169.* 22. Pests of Banana in Jamaica. *Jamaica Agri. Dept. Ent. Cir. No. 14.* (1934). 23. A Pasture Hand Book. *U. S. Agri. Dep. Misc. Pub. No. 194.* (1934). 24. Uniformity of Cotton Fiber Determined by Field Inspection. *U. S. Agri. Dep. Cir. No. 310.* (1934). 25. Feeding Care, and Management of Young Dairy Stock. *U. S. Agri. Dep. Farm. Bull. No. 1723.* (1934). 26. Cotton Breeding Investigations (1928--1932). *Mexico Agri. Exp. Stn. Bull. No. 217.* (1934). 27. Breeding for Egg Production. *Massachusetts Agri. Exp. Stn. Bull. No. 307.* (1934). 28. Silage Investigations. *Missouri Agri. Exp. Stn. Res. Bull. No. 22.* (1916). 29. Some factors favouring or opposing Fruitfulness in Apples. *Missouri Agri. Exp. Stn. Res. Bull. No. 32.* (1918). 30. The Normal Growth of Dairy Cattle. *Missouri Agri. Exp. Stn. Res. Bull.*

No. 36. (1920). 31. The effect of a varying moisture supply upon the development and composition of the maize plant at different periods of growth. *Missouri Agri. Exp. Stn. Res. Bull. No. 76.* (1925). 32. The effect of a varying supply of nutrients upon the character and composition of the maize plant at different periods of growth, *Missouri Agri. Exp. Stn. Res. Bull. No. 42.* (1921). 33. Physiology of Reproduction in Horticultural Plants; I. Reproduction and Metabolic Efficiency in the Tomato. *Missouri Agri. Exp. Stn. Res. Bull. No. 90.* (1926). 34. Physiology of Reproduction in Horticultural Plants; II. The Physiological Basis of Intermittent Sterility with Special Reference to the Spider Flower. *Missouri Agri. Exp. Stn. Res. Bull. No. 106.* (1927). 35. A Comparison of Four Systems of Pruning Grapes. *Missouri Agri. Exp. Stn. Res. Bull. No. 121.* (1928). 36. Social, Economic, and Homemaking Factors in Farm Living. *Missouri Agri. Exp. Stn. Res. Bull. No. 148.* (1930). 37. Taxonomy of the Genus *Phytophthora* de Bary. 38. The Anatomy of the Mammary Gland of Cattle. 39. Behaviour of Potassium and Sodium during the process of Soil Formation. 40. Cytological Observations of Deficiencies Involving Known Genes, Translocations and an Inversion in the *Zoomas*. *Missouri Agri. Exp. Stn. Res. Bull. Nos. 153, 160, 162, 163.* (1931) 41. The Influence of Systems of Cropping and Methods of Culture on Surface Runoff and Soil Erosion. 42. The effect of Ultra-Violet Rays on the Dermatitis Preventing Vitamin. *Missouri Agri. Exp. Stn. Res. Bull. Nos. 177, 178.* (1932). 43. The Distribution of the Genus *Phytophthora*. 44. Effects of Variations in the Amounts of Vitamin B and Protein in the Ration. 45. Power, Labor and Machine Costs in Crop Production. 46. The Nature of Shedding of Immature Apples. *Missouri Agri. Exp. Stn. Res. Bull. Nos. 184, 195, 197, 201.* (1933).

D. Circulars, Leaflets.

47. Lamb Dysentery. *Eng. Min. Agri. & Fish. Adv. Leaf. No. 214.* 48. Tuba. Root (Derris Spp.). 49. Gingelly (Sesamum Indicum). 50. Ground-nut or Peanut, (*Arachis Hypogaea*). 51. Bananas, (*Musa* Vars.). 52. Pepper (*Peper Nigrum*), S. S. & F. M. S. *Agri. Dep. Agri. Leaf. Nos. 1, 2, 4, 5, 9.*

E. Translations.

53. Vysotskii: New Methods of Cotton Cultivation, Hybridization, Vegetative Propagation, New Methods of Transplantation. 54. Moshkov; Photoperiodism in Trees and its Practical Importance. 55. Janini Janini: Artificial Fertilization of oranges, Production of New Varieties.

F. New Periodicals.

Foultry World. (Weekly). Feathered World. (Weekly).

ADDITIONS TO THE LIBRARY, JULY 1934.

A. Books.

1. Modern Coffee Planting *Windle, E. G.* (1934).
2. The Planning of Agriculture. *Astor (Viscount) & Murray, K. A. H.* (1933).
3. The Young man in Farming. *Getman, A. K. & Chapman, R. W.* (1933).
4. World Sugar Production and Consumption. *Robertson, C. J.* (1934).
5. Flowers and Gardens in India by Mrs. Temple Wright.—8th Edn. *Burns (W.); Rev.* (1934). Revised with a Chapter on Lawns and Rose Cultivation. *Davis, W. H.* (1934).
6. Pruning in South Africa. *Powell, H. C.* (1929).
7. Keeping and Using Farm Records. *Hopkins, J. A.* (1932).
8. Farmers' Practical Account Book. *Horrocks, W. A.* (1931).
9. Suppression of weeds by Fertilizers and Chemicals *Long, H. C.* (1934).
10. Practical Animal Husbandry. *Miller, M. C. & Robertson, E. D. S.* (1934).
11. Animal Management. *Creedy, H. J.* (1933).
12. Poultry Keeping in India—by Tweed—6th Edn. (Revised). *Mrs. Lord, M. V.* (1927).
13. Analytical Judgment Accounts for correct decisions of every description. *Dunlop, W. R.* (1933).
14. Analytical Judging Book (Supplement), *Dunlop, W. R.* (1934).
15. The Eradication of Bovine Tuberculosis. *Jordan, L.* (1933).
16. Hoare's Veterinary Materia Medica and Therapeutics—5th Edition—Revised. *Graig, J. R.* (1933).
17. Sewage Disposal in India and the East. *Williams, G. B.* (1924).
18. Internal Combustion Engines. *Maleev, V. L.* (1933).
19. Introduction to Bio-Chemistry. *Fearon, W. R.* (1934).
20. Fluorescence Analysis in Ultra-Violet Light. *Radley, J. A. & Grant J.* (1933).
21. Colloid Symposium Monograph, No. 10. *Weiser (H. B.) Ed.* (1932).
22. Virus Diseases of Plants. *Grainger, J.* (1934).
23. Researches on Fungi—Vol. 4 & 5. *Buller, A. H. R.* (1933).
24. Principles of Animal Biology—4th Edn.
25. Laboratory Directions in Principles of Animal Biology—4th Edn. *Shull, A. F. & Co* (1934).
26. Calculations and Interpretations of Analysis of Variance and Co-Variance.
27. Madras District Gazetteers—Ganjam District—Statistical Appendix, Volume 3.
28. Madras District Gazetteers—Kistna District—Statistical Appendix, Volume 2. *Snedecor, G. W.* (1934).

(Books for Students' Library.)

29. Company Accounts by A. Coles. 4th Edition. *Wood W. C.* (1930).
30. Students' Complete Book-Keeping—Accounting and Banking. 43rd Edn. *Fieldhouse, A.* (1933).
31. Advanced Accounts. *Carter, R. N.* (1933).
32. Partnership Law and Accounts. *Holland, R. W.* (1931).
33. Intermediate Book-Keeping, 5th Edn. *K. Subramania Ayyar.* (1930).
34. Book-Keeping and Accounts, 8th Edn. *Spicer, E. C. & Pegler, E. C.* (1931).
35. Higher Book-Keeping and Accounts by L. C. Cropper—5th Edn. (Revised). *Morris (F. D.), Rev. & Fison (A. K.), Rev.* (1931).

B. Annual Reports & Publications.

1. Review of the Sugar Industry of India during 1932—1933. *Indian Trade Journal Supplement, July, 1931.*
2. The Teachers' College Saidapet, Calendar for 1934—1935.
3. Annual Report of the Mysore Agricultural Department for 1932—1933.
4. Annual Report of the Imperial Bureau of Soil Science. England for 1933—1934.

C. Special Publications, Bulletins, Memoirs, &c.

5. Crop Planning Conference of Discussions. June 1934—Resume.
6. Bombay Weights and Measures Act of 1932.
7. Development of the Sugar Industry in Mysore. *Mys. Agri. Deptt. Bull. 18, 1931.*
8. A Preliminary Note on the leaf-cure Mosaic Disease of Sandal, by M. G. Venkata Rao, Bangalore. *Mysore Sandal Spike Invest. Comm. Bull. 3, 1933.*
9. Report on the Disease Situation in Coffee Areas in 1933.
10. Vegetable Oils as spreaders for Bordeaux Mixture. *Mys. Coffee Exp. Stn. Circular Nos. 7, 2, 1931.*
11. The Relation between exchangeable Sodium and Crop yield in Punjab soils and a new method of characterising Alkali Soils. *Punjab Irrigation Res. Pub. Vol. IV, No. 5, 1933.*
12. A Simple Method for determining the Reaction and Titration curves of Soils. *Punjab Irrigation Res. Pub. Vol. IV.*

No. 6, 1931. 13. Studies in the Cost of Production in the Punjab. 14. Economics of Cut Making in the Punjab. 15. Cost of Ginning and Pressing Cotton in the Punjab. 16. Economic Conditions of Simla Rickshaw Men. 17. Finance and Marketing of Cultivators' Wheat in the Punjab. *Board of Econ. Inquiry, Punjab. Pub. Nos. 33, 34, 36, 37, 38, 1934.* 18. "Dunlop" Pneumatic Tyres. Wheels and Hubs for Farm Carts. *Min. of Agri. & Fish. Lond. Cert. & Reports No. 46, 1933.* 19. The Residual Values of Feeding Stuffs and Fertilizers—Report by a Joint Committee of Scientific and Professional Men. *D. pt. of Agri. Scot. Misc. Pub. No. 7, 1932 Ldn.* 20. Tuberculous Infection in Milk. *Privy Council Med. Res. Spl. Report Ser. No. 189, 1933.* 21. Some Diseases of Farm Animals. 22. Asparagus. 23. Tomatoes: Cultivation, Diseases and Pests. 24. Fungus and Other Diseases of Crops (1928-32). *Min. of Agri. & Fish. Lond. Bull. Nos. 1, 60, 77, 79, 1934.* 25. Research in Agricultural Land Utilization—Scope and Method. *Social Sci. Res. Council. Bull. No. 2, June 1931.* 26. Research in Agricultural Income; Scope and Method. 27. Research in Agricultural Co-operation. *Social Sci. Res. Council. Bull. Nos. 6, 15, 1933.* 28. Experiments with Forage Crops at the Fort Hays Branch Station, Hays, Kans. (1913-1928). *U. S. Agri. Dept. Tech. Bull. No. 410, April, 1934.* 29. The Nitrogen Nutrition of Tobacco. *U. S. Agri. Dept. Tech. Bull. No. 414, 1934.* 30. Greenhouse Construction and Heating. *U. S. Agri. Farm. Bull. 1318, 1934.* 31. Hilgardia.—The penetration of Insecticidal Oils into Porous Solids. *Calif. Agri. Exp. Stn. Hilgardia, Vol. 8, No. 2, Nov. 1933.* 32. —Sulphuric Acid as a penetrating Agent in Arsenical Sprays for Weed Control. *Calif. Agri. Exp. Stn. Hilgardia, Vol. 8, No. 4, Dec. 1933.* 33. —Principles Governing the Reclamation of Alkali Soils. *Calif. Agri. Exp. Stn. Hilgardia, Vol. 8, No. 5, Jan. 1934.* 34. The Value of Sprays and Fumigation for Resistant Black Scale Control. *Calif. Fruit. Grow. Exchange, (Lib. Acc. No. 30862), July, 1925.* 35. Hand Book of Citrus Insect Control for 1926. *Calif. Fruit. Grow. Exchange, Bull. No. 3, July, 1926.* 36. Hand Book of Citrus Insect Control for 1927. *Calif. Fruit. Grow. Exchange, Bull. No. 4, July, 1927.* 37. Hand Book of Citrus Insect Control for 1928. *Calif. Fruit. Grow. Exchange, Bull. No. 5, July, 1928.* 38. —The Mediterranean Fruit Fly. *Calif. Fruit. Grow. Exchange, Bull. No. 6, July, 1929.* 39. Hand Book of Citrus Insect Control for 1930. *Calif. Fruit. Grow. Exchange, Bull. No. 7, July, 1930.* 40. Hand Book of Citrus Insect Control for 1931. *Calif. Fruit. Grow. Exchange, Bull. No. 8, July, 1931.* 41. Hand Book of Citrus Insect Control for 1932. *Calif. Fruit. Grow. Exchange, Bull. No. 9, July, 1932.* 42. Hand Book of Citrus Insect Control for 1933. *Calif. Fruit. Grow. Exchange, Bull. No. 10, July, 1933.* 43. Wild Rice of Kwangtung and New Variety Bred from the Hybrids of Wild Rice with Cultivated Rice. *Rice Exp. Stn. Canton, China, Pub. Ser. No. III, Agron. Bull. 3, December, 1933.* 44. The Sterile Flower of Wild Rice of Kwangtung. *Rice Exp. Stn. Canton, China, Pub. Ser. No. III, Agron. Bull. 4, March, 1934.*

D. Circulars, Leaflets, &c.

45. Cabbage Butterfly Caterpillars. 46. Cabbage Moth Caterpillars. 47. The Barn Owl Caterpillars. 48. Nest Boxes for Birds Caterpillars. *Min. of Agri. & Fish. Lond. Adv. Leaf. Nos. 69, 70, 211, 212, 1934.*

E. Translations (of the Imperial Bureau of Plant Genetics, England):

49. Directions for Making Square Yard Harvests of Legumes, by C. J. Willard. 50. Proceedings for Making Draft Tests of Plows and the Use of Soil Sampling Tube. 51. Research in Mechanical Phases of Cotton Ginning, by Bennet and Some Mechanical Elements involved in Good Ginning, by Bennet. *U. S. Agri. Deptt. Bur. of Agri. Engineering.* 52. Water Melons of the Northern Hemisphere, by Pangalo (K. I.):

F. Reprints.

53. Rice Research in Madras, by K. Ramiah Reprinted from "Current Science," Vol. III, No. 1, July, 1934 (pp. 34-36).

ADDITIONS TO THE LIBRARY, AUGUST 1934

A. Books.

1. Studies in Sugar Boiling translated by O. W. Willcox. *J. G. Thieme*. (1928).
2. Research in Agricultural Credit. *Social Science Research Council Pubn.* (1931).
3. Research in Farm Management. 4. Research in Agricultural Insurance. *Social Science Research Pubn.* (1932).
5. Research in Farm Labour. 6. Research in Farm Real Estate Values. 7. Research in Agricultural Land Tenure. 8. Research in Agricultural Policy. *Social Science Research Council Pubn.* (1933).
9. The Asylum Press Almanac and Directory—1934. *Social Science Research Council Pubn.* (1934).
10. History of Services of Gazetted and other officers in the Civil Department serving in the Madras Presidency—corrected upto July 1934. (1934).
11. Monthly Rainfall of India—1931. (1934).
12. Coimbatore Agricultural College Calendar,(1934-'35). (1934).

B. Reports.

1. Cochin Agricultural Department Administration Report for 1932-'33. 2. Annual Report (twenty-first year) of the East Malling Research Station for 1933. 3. Forty-third Annual Report (Fiscal year ending June 30, 1932) of the Agricultural Experiment Station of the Alabama Polytechnic Institute, Auburn. 4. Fifty-sixth Report (with bulletins) of the Connecticut Agricultural Experiment Station, New Haven, for the year 1932.

C. Bulletins, Memoirs, &c.

5. Notes on Horticulture—Fruits by R. P. Misra. (1934).
6. Notes on an Enquiry into certain Public Health Aspects of village life in India and population and health in India—The Real Problem—by Major-General Sir John Megaw, K.C.I.E., K.H.P., I.M.S., Director-General, Indian Medical Service. (1934).
7. Bibliography of Publications Relating to Coffee Soils and Fertilisers. *Imp. Bur. of Soil Science 1934, Publications.* (1934).
8. The Scientific Principles of Poultry Feeding. 9. Specifications and Methods of Analysis for certain Insecticides and Fungicides. *Eng. Min. Agri. & Fisheries, Bull. Nos. 7, 82, 1934.*
10. Fodders and Feeding Stuff in Malaya by C. D. V. George, O. B. E., &c. *Agri. Dept. S. S. & F. M. S. General Ser. No. 17, 1934.*
11. Citrus Wastage Investigations, Progress Report No. 2—Season 1933 by V. A. Putterill.
12. Some Important Aspects in the Manufacture and Marketing of Dairy Products by H. B. Davel. *Agri. Dept. U. S. Africa. Bull. No. 131, 135, 1934.*
13. A Simple System of Farm Accounts and Records by Liversage.
14. Preliminary Observations on Coffee Roots in Kenya by A. D. Trench.
15. Report of Committee on Coffee Berry Disease. *Konya Agri. Dept. Bull. Nos. 1, 2, 3, 1934.*
16. The Rices of Northern Nyasaland by Frank Barker. *Nyasaland Agri. Dept. Bull. 12, 1934.*
17. Report on the Study of the Irrigation of Water of Rice. *Sun Yatson Univer. Agri. Coll. Bull. 1 of 1929.*
18. Preliminary Report on the Fertilizer Experiments of Rice. *Sun Yatson Univer. Agri. Coll. Bull. 2, 1932.*
19. One Year of the AAA; the Record Reviewed. *U. S. Agri. Dept. Adjust. Admn. June Issuc. 1934.*
20. Brucellosis—A Public Health Problem by Ward Giltner. *Mich. Agri. Exp. Stn. Memr. 1, March 1934.*
21. Egg Weight in the Domestic Fowl by E. M. Funk & H. L. Kempster. *Missouri Agri. Exp. Stn. Bull. 332, 1934.*
22. Chromosome Structure in *Tradescantia* V. Optical Analysis of a Somatic Telophase Chromosome by B. R. Nebel. *New York State Agri. Exp. Stn. Tech. Bull. 220, 1933.*
23. Crown Gall and Hairy Root of Apples in Nursery and Orchard—by W. O. Gower. *New York State Agri. Exp. Stn. Tech. Bull. 638, 1934.*
24. The Effect of Time and Rate of Application of Nitrate of Soda on the Yield of Cotton. *Texas Agri. Exp. Stn. Bull. 490, 1934.*
25. The Influence of Some

Factors on the Hatchability of the Hens Egg. 26. Varieties of Sorghum in Kansas. *Kansas Agri. Exp. Stn. Tech. Bull. Nos. 37, 266, 1934.* 27. Root Stock Effects with Cherries—Seed & Phyton Propagation. 28. Fixation and Penetration of Phosphates in Vermont Soils by V. L. Weiser. 29. Land Utilization as a Basis of Rural Economic Organisation—based on a Study of Land Utilization and related problems in 13 Hill Towns of Vermont. 30. The Carbohydrate Contents of the Maple Tree. 30(a). The Feeding Value of Artificially Dried Young Grass—II. 31. Forty-sixth Annual Report (1932-'33)—by J. L. Hills. *Vermont Agri. Exp. Stn. Bull. Nos. 352, 356, 357, 358, 359, 360, 1933.* 32. Manganese—An Essential Element for Green Plants. 33. The Relative Growth and Development of Corn Varieties of Widely different Maturity Dates During Successive Time Intervals throughout their Life Cycle—R. G. Wiggans. 34. Longevity of Rhizobium Japonicum in Relation to its Symbiont on the Soil—J. K. Wilson. *Cornell Univ. Agri. Exp. Stn. Mem. Nos. 151, 152, 162, 1934.* 35. Cost Accounts on New York Farms. *Cornell Agri. Univ. Exp. Stn. Bull. No. 554, April 1933.* 36. Studies in Plant Propagation. *Cornell Agri. Univ. Exp. Stn. Bull. No. 571, June 1933.* 37. The Rate of Photosynthesis of Apple Leaves under Natural Conditions—Part I. *Cornell Agri. Univ. Exp. Stn. Bull. No. 577, Novr. 1933.* 38. Pruning and Training Tomatoes. *Cornell Agri. Univ. Exp. Stn. Bull. No. 580, Juny. 1934.* 39. Factors Influencing the Occurrence of Potato Scab in New York. 40. An Economic Study of the Marketing of Certain Perishable Farm Products in Albany, New York. *Cornell Agri. Univ. Exp. Stn. Bull. Nos. 581, 585, Feby. 1934.* 41. A Study of the Effect of Removing Foremilk on the Fat Content of the Remainder of the Milking. *Cornell Agri. Univ. Exp. Stn. Bull. No. 589, Mar. 1934.* 42. Production of Early Blooms of Chrysanthemums by the Use of Black Cloth to Reduce the Length of Day. *Cornell Agri. Univ. Exp. Stn. Bull. No. 594, Apr. 1934.*

D. Circulars, Leaflets, &c.

43. Contagious Diseases of Poultry and their treatment. 44. Non-Contagious Diseases of Poultry and their treatment. 45. Common Affections of Poultry and their treatment. *Madras Vetr. Dept. Leaf. Nos. XIII, XIV, XV, 1934.* 46. Improved Varieties of Rice for Sind. *Sind Agri. Dept. Leaf. No. 3 (1st Edn. Jan. 1933).* 47. Improved Varieties of Jowar for Sind. *Sind Agri. Dept. Leaf. No. 4 (1st Edn. Apr. 1931).* 48. Useful Agricultural Implements for Sind. 49. Insect Pests on Crops in Sind. *Sind. Agri. Dept. Leaf. Nos. 5, 6 (1st Edn. July 1931).* 50. Scum on the Water Surface of Rice Fields. *Sind Agri. Dept. Leaf. No. 7 (1st Edn. Novr. 1931).* 51. Agricultural Propaganda by Touring Parties. 52. Horticultural Series—Propagation of Plants—by Seed and by Cuttings. 53. Horticultural Series—Propagation of Plants—by Layering. *Sind. Agri. Dept. Leaf. Nos. 8, 9, 10 (1st Edn. Jan. 1932).* 54. Horticultural Series—Propagation of Plants—by Grafting. 55. Horticultural Series—Propagation of Plants—by Budding. 56. The Cultivation and Use of Berseem in Sind. *Sind. Agri. Dept. Leaf. Nos. 11, 12, 13 (1st Edn. Feby. 1932).* 57. Green Manuring. *Sind. Agri. Dept. Leaf. No. 14 (1st Edn. Aug. 1932).* 58. Potato Cultivation in Sind. 59. Groundnut Cultivation in Sind. 60. Useful Agricultural Implements for Sind—The Egyptian ("SIRCAR") Plough. 61. Onion Cultivation in Sind. *Sind. Agri. Dept. Leaf. Nos. 15, 16, 17, 18 (1st Edn. Novr. 1932).* 62. Green Manuring of Cotton with Guar. *Sind. Agri. Dept. Leaf. No. 19 (1st Edn. Jany. 1933).* 63. Horticultural Series—The Establishment of a Fruit Tree Orchard. 64. Useful Agricultural Implements for Sind—"COOPER" No. 25 (Meston Type) Plough. *Sind. Agri. Dept. Leaf. Nos. 20, 21 (1st Edn. Feby. 1933).* 65. Control of Green Smut ('Kani') on Jowar. *Sind. Agri. Dept. Leaf. No. 22 (1st Edn. Mar. 1933).* 66. Calendar of Main Agricultural Operations at the Agricultural Research Station, Sakrand. *Sind. Agri. Dept. Leaf. No. 23 (1st Edn. Feby. 1933).* 67. The Agricultural Department in Sind. *Sind. Agri. Dept. Leaf. No. 24 (2nd Edn. Apr. 1933 & 1st Edn. Leaf. 1, 1931).* 68. The Preparation of Lands,

Formerly under Rice Cultivation, to render them suitable for dry crop cultivation. 69. The Cultivation of Oats as a Green Fodder in the Rabi Season. 70. Castor Cultivation in Sind. *Sind. Agri. Dept. Leaf. Nos. 25, 26, 27* (1st Edn. May 1933). 71. Horticultural Series—The Protection of Fruits from Frost. *Sind. Agri. Dept. Leaf. No. 28* (1st Edn. June 1933). 72. The Rat Pest on Rice & Wheat. *Sind. Agri. Dept. Leaf. No. 29* (1st Edn. May 1933). 73. The Cultivation of Rabi Oil-Seed in the Barrage Areas of Sind. *Sind. Agri. Dept. Leaf. No. 30* (1st Edn. Aug. 1933). 74. Cotton Cultivation on the Indus Right Bank Areas. *Sind. Agri. Dept. Leaf. No. 31* (1st Edn. Novr. 1933). 75. Cultivation of Kangni Rice in Karachi District. *Sind. Agri. Dept. Leaf. No. 32* (1st Edn. Decr. 1933). 76. Useful Agri. Implements for Sind—Three Coultured Rabi Seed-Drill. *Sind. Agri. Dept. Leaf. No. 33* (1st Edn. Jany. 1934). 77. The Sowing of the Cotton Crop by Drill method. *Sind. Agri. Dept. Leaf. No. 34* (1st Edn. Feby. 1934). 78. The Irrigation of Dry Crops in the Barrage Areas including a Suggested-time of Agricultural Operations and Distribution of Irrigation Water. 79. The Cultivation of Kharif Oil-Seed Crops in the Barrage Areas of Sind. *Sind. Agri. Dept. Leaf. Nos. 35, 36* (1st Edn. Mar. 1934). 80. Improved Varieties of Jowar for Sind. *Sind. Agri. Dept. Leaf. No. 37* (2nd Edn. May 1934). 81. Tomato Culture in Missouri. *Missouri Agri. Exp. Stn. Cir. No. 173*, Feby, 1934. 82. The Soybean Crop in Missouri. *Missouri Agri. Exp. Stn. Cir. No. 174*, Mar. 1934. 83. Liming Wisconsin Soils. *Univ. Wis. Agri. Cir. 266*, Feby. 1934. 84. The Home Garden & Orchard in the Wichita Valley. *Texas Agri. Exp. Stn. Cir. No. 70*, Jany. 1934.

E. Special Periodicals, Publications & Reprints.

85. A Gravitational Survey of the Sub-Alluvium of the Jhelum-Chenab Ravi Doabs, and its Application to Problems of Waterlogging. *Punjab Irr. Res. Instt. Res. Pubn. Vol. VI, No. 1*, Feby. 1934. 86. Outline of Results from the Study of the Irrigation Water of Rice at Canton. *Lingnan Science Jrl. Vol. 9, No. 3 (Canton)*, 1930.

F. Translations (By I. A. B.)

87. Some Experiments on the relation and the profitableness of different varieties to increased nitrogenous manuring. (*Z. Pflanzenenahrung, Dungung u. Bodenkunde* 1931; 10; 169-81).

G. New Periodicals

1. Proceedidgs of the Indian Academy of Sciences—Irregular. 2. Harper Adams Agricultural College Poultry Utility Journal—Monthly. 3. "Eggs"—Weekly.

ADDITIONS TO LIBRARY, SEPTEMBER 1934.

A. Books.

1. Weather.—Rev. Edn. *Abercromby, R. & Goldie, A. H. R.* (1934)
2. Rural Refuse and its Disposal. *Ashford, E. B. & Baker, H.* (1934).
3. A Text Book of West African Agriculture. *Irvine, F. R.* (1934).
4. The Apple. *Hall, A. D. & Crane, M. B.* (1933).
5. Modern Flowergrowing for Profit. *Shawell-Cooper, W. C.* (1933).
6. The Wright Encyclopaedia of Gardening. *Wright, W. P.* (1933).
7. Farm Accounts in the Punjab,—(1932-'33). *Karter Singh & Arjan Singh.* (1934).
8. The Foundations of Agricultural Economics—2nd Edn. (Rev. & Enlarged). *Venn, J. A.* (1933).
9. The Principles and Practice of Marketing. *Simmat, R.* (1933).
10. Veterinary Hygiene,—2nd Edn. Rev. *Linton, R. G.* (1934).
11. Poultry Diseases. *Kaup, B. F.* (1933).
12. Recent Advances in Plant Physiology, 2nd Edn. *Barton-Wright, E. C.* (1933).
13. Selective Fertilization. *Jones, D. F.* (1928).
14. Taxonomy of the Flowering Plants. *Johnson, A. M.* (1931).
15. Text Book of Systematic Botany, 2nd Edn Rev. *Swingle, D. B.* (1934).
16. Introduction to Cytology, 3rd Edn. Rev. *Sharp, L. W.* (1934).
17. Virus Diseases of Plants: A Bibliography. (1934).
18. The Vitamins, 2nd Edn. Rev. *Sherman, H. C. & Smith, S. L.* (1934).
19. Bio-Chemistry of Nitrogen, Conservation. *Fowler, G. F.* (1934).
20. The Atom. *Tutin, J.* (1934)
21. Perkin and Kippings Organic Chemistry, Part III. *Kipping, F. S. & Kipping, F. B.* (1934).
22. A Comprehensive Treatise on Inorganic and Theoretical Chemistry, Vol. 13—Iron (Part 2). *Mellor, J. W.* (1934).
23. Colloid Symposium Annual, Vol. 7. *Weiser, H. B. (Ed.)* (1930).
24. Inorganic Colloid Chemistry, Vol. 1, the Colloidal Elements. *Weiser, H. B.* (1933).
25. Laboratory Manual of Colloid Chemistry, 3rd Edn Rev. *Holmes, H. N.* (1934).
26. A General Text Book of Entomology, 3rd Edn. Rev. *Inms, A. D.* (1934).
27. Morphology of the Insect Abdomen, Part I—General Structure of the Abdomen and its appendages. *Snodgrass, R. E.* (1931).
28. Morphology of the Insect Abdomen. Part II the genital Ducts and the Ovipositor. *Snodgrass, R. E.* (1933).
29. The Life of the Bee, translated by A. Sutro, (Reprint Edition). *M. Maerlueck* (1930).
30. The Way of the Bee, translated by P. Kirwan. *Rendl, G.* (1933).
31. The Progress of Science. *Crowther, J. G.* (1934).
32. The New Background of Science; 2nd Edition. *Jean, J.* (1934).

B. Reports.

1. Report of the Botanical Survey of India for 1931-'32.
2. Nyasaland Protectorate Agri. Deptt.—Annual Report for 1931-'32.
3. Tanganyika Agricultural Department,—Annual Report for 1933.
4. Madras Veterinary College Calendar for 1934-'35.
5. Madras Engineering College Calendar for 1934-'35.

C. Special Publications, Memoirs, &c.

6. Market Practices in the Punjab *Punjab Board of Econ. Enq. Pubn.* 39. (1934).
7. Entomological Investigations on the Spike Diseases of Sandal, Jassidae. *Cicindelidae (Col.)* and supplementary data on Neuroptera and Elateridae (Col.).
8. *Coccinellidae (Col.)*.
9. *Coccinellidae (Col.)*.
10. *Coccinellidae (Col.)*.
11. *Fulgoridae (Homopt.)*
12. The spike disease of Sandal.—On the life-history and Morphology of *Petaloccephala Nigrilinea Walk.*, (Jassidae, Homopt.) *Indian Forest Records, Vol. 19, Nos. 4, 5, 6, 7, 8, 9.* (1934).
13. Entomological Investigations Studies on Insect Transmission.
14. Immature stages of Indian Coleoptera (14).
15. On the Biology of Mantidae (Orthopt.)
16. The spike disease of sandal (21).
17. Formicidae (Hymen).
18. Anthicidae (Col.).
19. Methods of testing the susceptibility of timbers to termite attack.
20. Immature stages of Indian Coleoptera (15) (Scolytidae).
21. Entomological Investigations on the spike disease of sandal (24) Pentatomidae. *Indian Forest Records, Vol. 26; *Parts 1, 2, 3, 4, 5, 6, 7, 8, 9.* (1934).
22. An Act to provide for the imposition and collection of an excise duty on sugar.
23. An Act to regulate the price of sugar-cane intended for use in sugar factories. *India Act Nos. 14, 15 of 1934.*
24. The Pea-nut; its history; Histology, Physiology and Utility by R. A. Waldron. Production of New cell formation in Plants by W. R. Taylor. Water content and Transpiration of leaves by A. W. Clark. Bio-Chemical Studies of Insectivorous Plants. *Pennsylvania University*

Botanical Laboratory Contributions, Vol. 4, No. 2. (1919). 25. A Statement of Objectives for Agriculture. *Iowa Agri. Deptt. Pubn.* (1933). 26. The Edible Fat Problem in Germany. *U. S. A. Bur. Agri. Econ. F. A. S. No. 61.* (1934). 27. German Animal Fat Protection Program. *U. S. A. Bur. Agri. F. A. S. "Foreign Markets"* Reprint, July, 1933. 28. Bread grain consumption and Trade in Scandinavian Countries. 29. Wheat consumption and Trade in Netherlands. *U. S. A. Bur. Agri. F. S. No. 60.* (1933). 30. Foreign Government Legislation affecting Wheat. *U. S. A. Bur. Agri. F. S. No. 58.* (1932). 31. Soviet Agrarian Policy. (1933). 32. The World Situation in Feed Grains (1934). 33. The World Sugar Situation. (July, 1934). 34. Cotton Production in the Anglo-Egyptian Sudan. *U. S. A. Bur. Agri. F. S. No. 62.* (1934).

D. Bulletins.

35. Irwin Canal Soil Survey Report Series I—Introductory and General Part. *Mys. Agri. Deptt. Pubn.* (1934). 36. The Determination of Exchangeable bases in Soil. *Eng. Imp. Bur. Soil Sci. Tech. Commu. No. 30.* (1934). 37. Grassland and Forage Crops in Thuringia, Czechoslovakia and Hungary. *Eng. Imp. Bur. Plant Gen. Herbage Pubn. Ser. Bull. 15.* (1934). 38. Turkeys. 39. Arable Crops of the Far n. *Eng. Min. Agri. Fish. Bull. Nos. 67, 72.* (1934). 40. Further Experiments on the inheritance of Chlorophyll Deficiency in New World Cottons. *Trinidad Cotton Res. Stn. Memoirs Ser. A—Genetics, No. 8.* (1934). 41. Raw Linseed Oil and Seal Oil for Controlling Irregular Blossoming and Foliation in Fruit Trees. *Union of South Africa, Agri. Deptt. Bull. No. 125.* (1934). 42. The External Anatomy of the Red Date Scale *Phoenicococcus Marlatti* Cockerell, and its Allies. 43. Forage Crop Field Experiments at West Point. Miss. 44. Fluctuations in Prices of Cotton Futures Contracts. 45. Influence of Storage Temperatures on the Rest Period and Dormancy of Potatoes. 46. Marketing Cantaloups and other Muskmelons. *U. S. A. Agri. Deptt. Tech. Bull. Nos. 404, 419, 423, 424, 425.* (1934). 47. The Relation of Air Conditions to Tobacco Curing. *Wisconsin Agri. Exp. Stn. Bull. No. 110.* July, 1931. 48. Physiological Dropping of Fruits—II. In Relation to Genetic Relationship of Plants. *Delaware Agri. Exp. Stn. Bull. No. 157.* Nov. 1931. 49. Wheat and Wheat By-Products for Laying Hens. *South Dakota Agri. Exp. Stn. Bull. No. 284.* May, 1934. 50. Comparative Metabolism of Several Calcareous Materials used in Poultry Feeding. *South Dakota Agri. Exp. Stn. Bull. No. 287.* Jan. 1934. 51. Relations between Orchard Soils and Cover Crops. *New York Agri. Exp. Stn. Bull. No. 632.* July, 1933.

E. Circulars.

52. Handling, Planting and Early Care of Fruit Trees. *Missouri Agri. Extn. Ser. Cir. 273.* (1928). 53. Pruning the Grape. 54. Establishment and Management of Vineyard. *Missouri Agri. Extn. Ser. Cir. Nos. 235, 236.* (1929). 55. Growing Barley for Malt and Feed. 56. Planning a Subsistence Homestead. *U. S. Agri. Deptt. Farm Bull. Nos. 1732, 1733.* (1934). 57. Traps for the Japanese Beetle and How to use them. *U. S. Agri. Deptt. Farm. Misc. Publ. No. 201.* (1934). 58. Protection of Orchard and Shade Trees and Ornamental Shrubs from Injury by the Japanese Beetle. *U. S. Agri. Deptt. Forul. Cir. No. 317.* July 1934. 59. Directions for Collecting and Preserving Insects. *Canada Agri. Deptt. Pamphlet 14.* Decr. 1931. 60. Cultivation of the Vegetable Marrow. 61. Seakale Cultivation. 62. Dry Rot of Potatoes. 63. Grain Weevils. 64. Barley Growing. *Eng. Min. Agri. Fish. Adv. Leaf. Nos. 216, 217, 218, 219, 220.* (1934).

F. Imperial Agricultural Bureau of Plant Genetics (Translations).

65. A Study on Millet by S. Lewicki. *Translations from Polish.* 66. Genetics at the service of Socialistic Agriculture by N. L. Vavilov. 67. The Principal Changes introduced by the Revolution into the Diversity of the cultivated plants by Shukovsky P. M. 68. Theoretical Basis of Plant Acclimatization, by Maleyev V. P. *Abridged Translation.*

G. Periodicals.

1. Poultry Tribune.—Monthly.